

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/21, 5/10, 15/18, 15/63, C12Q 1/00, 1/68, 1/70, A61K 38/18, C07K 14/50, 16/22	A1	(11) International Publication Number: WO 99/46364 (43) International Publication Date: 16 September 1999 (16.09.99)
(21) International Application Number: PCT/US99/05021 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 09/042,105 13 March 1998 (13.03.98) US 09/107,997 30 June 1998 (30.06.98) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). CAO, Liang [CN/US]; Flat 32D, Block 13A, South Horizons, Hong Kong (CN). HU, Jing-Shan [CN/US]; 1247 Lakeside Drive #3034, Sunnyvale, CA 94086 (US). (74) Agents: WALES, Michele, M. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With an indication in relation to deposited biological</i> <i>material furnished under Rule 13bis separately from the</i> <i>description.</i>	
(54) Title: VASCULAR ENDOTHELIAL GROWTH FACTOR 2 (57) Abstract Disclosed are human VEGF-2 polypeptides, biologically active, diagnostically or therapeutically useful fragments, analogs, or derivatives thereof, and DNA(RNA) encoding such VEGF-2 polypeptides. Also provided are procedures for producing such polypeptides by recombinant techniques and antibodies and antagonists against such polypeptides. Such polypeptides and polynucleotides may be used therapeutically for stimulating wound healing and for vascular tissue repair. Also provided are methods of using the antibodies and antagonists to inhibit tumor angiogenesis and thus tumor growth, inflammation, diabetic retinopathy, rheumatoid arthritis, and psoriasis.		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/21, 5/10, 15/18; 15/63, C12Q 1/00, 1/68, 1/70, A61K 38/18, C07K 14/50, 16/22	A1	(11) International Publication Number: WO 99/46364 (43) International Publication Date: 16 September 1999 (16.09.99)
(21) International Application Number: PCT/US99/05021 (22) International Filing Date: 10 March 1999 (10.03.99) (30) Priority Data: 09/042,105 13 March 1998 (13.03.98) US 09/107,997 30 June 1998 (30.06.98) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). CAO, Liang [CN/US]; Flat 32D, Block 13A, South Horizons, Hong Kong (CN). HU, Jing-Shan [CN/US]; 1247 Lakeside Drive #3034, Sunnyvale, CA 94086 (US). (74) Agents: WALES, Michele, M. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With an indication in relation to deposited biological</i> <i>material furnished under Rule 13bis separately from the</i> <i>description.</i>	
(54) Title: VASCULAR ENDOTHELIAL GROWTH FACTOR 2 (57) Abstract Disclosed are human VEGF-2 polypeptides, biologically active, diagnostically or therapeutically useful fragments, analogs, or derivatives thereof, and DNA(RNA) encoding such VEGF-2 polypeptides. Also provided are procedures for producing such polypeptides by recombinant techniques and antibodies and antagonists against such polypeptides. Such polypeptides and polynucleotides may be used therapeutically for stimulating wound healing and for vascular tissue repair. Also provided are methods of using the antibodies and antagonists to inhibit tumor angiogenesis and thus tumor growth, inflammation, diabetic retinopathy, rheumatoid arthritis, and psoriasis.		

VASCULAR ENDOTHELIAL GROWTH FACTOR 2

Background of the Invention

5 The present invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production of such polynucleotides and polypeptides. The polypeptides of the present invention have been identified as members of the vascular endothelial growth factor family. More particularly, the polypeptides of the present invention are
10 human vascular endothelial growth factor 2 (VEGF-2). The invention also relates to inhibiting the action of such polypeptides.

The formation of new blood vessels, or angiogenesis, is essential for embryonic development, subsequent growth, and tissue repair. Angiogenesis is also an essential part of certain pathological conditions, such as neoplasia (*i.e.*, tumors and
15 gliomas). Abnormal angiogenesis is associated with other diseases such as inflammation, rheumatoid arthritis, psoriasis, and diabetic retinopathy (Folkman, J. and Klagsbrun, M., *Science* 235:442-447(1987)).

Both acidic and basic fibroblast growth factor molecules are mitogens for endothelial cells and other cell types. Angiotropin and angiogenin can induce
20 angiogenesis, although their functions are unclear (Folkman, J., *Cancer Medicine*, Lea and Febiger Press, pp. 153-170 (1993)). A highly selective mitogen for vascular endothelial cells is vascular endothelial growth factor or VEGF (Ferrara, N. *et al.*, *Endocr. Rev.* 13:19-32 (1992)), which is also known as vascular permeability factor (VPF).

25 Vascular endothelial growth factor is a secreted angiogenic mitogen whose target cell specificity appears to be restricted to vascular endothelial cells. The murine VEGF gene has been characterized and its expression pattern in embryogenesis has been analyzed. A persistent expression of VEGF was observed in epithelial cells adjacent to fenestrated endothelium, *e.g.*, in choroid plexus and kidney glomeruli.
30 The data was consistent with a role of VEGF as a multifunctional regulator of endothelial cell growth and differentiation (Breier, G. *et al.*, *Development* 114:521-532 (1992)).

VEGF shares sequence homology with human platelet-derived growth factors, PDGF α and PDGF β (Leung, D.W., *et al.*, *Science* 246:1306-1309, (1989)). The
35 extent of homology is about 21% and 23%, respectively. Eight cysteine residues contributing to disulfide-bond formation are strictly conserved in these proteins.

Although they are similar, there are specific differences between VEGF and PDGF. While PDGF is a major growth factor for connective tissue, VEGF is highly specific for endothelial cells. Alternatively spliced mRNAs have been identified for both VEGF, PLGF, and PDGF and these different splicing products differ in biological activity and in receptor-binding specificity. VEGF and PDGF function as homo-dimers or hetero-dimers and bind to receptors which elicit intrinsic tyrosine kinase activity following receptor dimerization.

VEGF has four different forms of 121, 165, 189 and 206 amino acids due to alternative splicing. VEGF121 and VEGF165 are soluble and are capable of promoting angiogenesis, whereas VEGF189 and VEGF-206 are bound to heparin containing proteoglycans in the cell surface. The temporal and spatial expression of VEGF has been correlated with physiological proliferation of the blood vessels (Gajdusek, C.M., and Carbon, S.J., *Cell Physiol.* 139:570-579 (1989); McNeil, P.L., *et al.*, *J. Cell. Biol.* 109:811-822 (1989)). Its high affinity binding sites are localized only on endothelial cells in tissue sections (Jakeman, L.B., *et al.*, *Clin. Invest.* 89:244-253 (1989)). The factor can be isolated from pituitary cells and several tumor cell lines, and has been implicated in some human gliomas (Plate, K.H., *Nature* 359:845-848 (1992)). Interestingly, expression of VEGF121 or VEGF165 confers on Chinese hamster ovary cells the ability to form tumors in nude mice (Ferrara, N. *et al.*, *J. Clin. Invest.* 91:160-170 (1993)). The inhibition of VEGF function by anti-VEGF monoclonal antibodies was shown to inhibit tumor growth in immune-deficient mice (Kim, K.J., *Nature* 362:841-844 (1993)). Further, a dominant-negative mutant of the VEGF receptor has been shown to inhibit growth of glioblastomas in mice.

Vascular permeability factor (VPF) has also been found to be responsible for persistent microvascular hyperpermeability to plasma proteins even after the cessation of injury, which is a characteristic feature of normal wound healing. This suggests that VPF is an important factor in wound healing. Brown, L.F. *et al.*, *J. Exp. Med.* 176:1375-1379 (1992).

The expression of VEGF is high in vascularized tissues, (*e.g.*, lung, heart, placenta and solid tumors) and correlates with angiogenesis both temporally and spatially. VEGF has also been shown to induce angiogenesis *in vivo*. Since angiogenesis is essential for the repair of normal tissues, especially vascular tissues, VEGF has been proposed for use in promoting vascular tissue repair (*e.g.*, in atherosclerosis).

U.S. Patent No. 5,073,492, issued December 17, 1991 to Chen *et al.*, discloses a method for synergistically enhancing endothelial cell growth in an appropriate environment which comprises adding to the environment, VEGF, effectors and serum-derived factor. Also, vascular endothelial cell growth factor C sub-unit DNA has been prepared by polymerase chain reaction techniques. The DNA
5 encodes a protein that may exist as either a heterodimer or homodimer. The protein is a mammalian vascular endothelial cell mitogen and, as such, is useful for the promotion of vascular development and repair, as disclosed in European Patent Application No. 92302750.2, published September 30, 1992.

10

Summary of the Invention

The polypeptides of the present invention have been putatively identified as a novel vascular endothelial growth factor based on amino acid sequence homology to
15 human VEGF.

In accordance with one aspect of the present invention, there are provided novel mature polypeptides, as well as biologically active and diagnostically or therapeutically useful fragments, analogs, and derivatives thereof. The polypeptides of the present invention are of human origin.

20 In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules comprising polynucleotides encoding full length or truncated VEGF-2 polypeptides having the amino acid sequences shown in SEQ ID NOS:2 or 4, respectively, or the amino acid sequences encoded by the cDNA clones deposited in bacterial hosts as ATCC Deposit Number 97149 on May 12, 1995 or
25 ATCC Deposit Number 75698 on March 4, 1994.

The present invention also relates to biologically active and diagnostically or therapeutically useful fragments, analogs, and derivatives of VEGF-2.

In accordance with still another aspect of the present invention, there are provided processes for producing such polypeptides by recombinant techniques
30 comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing a nucleic acid sequence encoding a polypeptide of the present invention, under conditions promoting expression of said proteins and subsequent recovery of said proteins.

In accordance with yet a further aspect of the present invention, there are
35 provided processes for utilizing such polypeptides, or polynucleotides encoding such polypeptides for therapeutic purposes, for example, to stimulate angiogenesis,

wound-healing, growth of damaged bone and tissue, and to promote vascular tissue repair. In particular, there are provided processes for utilizing such polypeptides, or polynucleotides encoding such polypeptides, for treatment of peripheral artery disease, such as critical limb ischemia and coronary disease.

5 In accordance with yet another aspect of the present invention, there are provided antibodies against such polypeptides and processes for producing such polypeptides.

In accordance with yet another aspect of the present invention, there are provided antagonists to such polypeptides, which may be used to inhibit the action of
10 such polypeptides, for example, to prevent tumor angiogenesis and thus inhibit the growth of tumors, to treat diabetic retinopathy, inflammation, rheumatoid arthritis and psoriasis.

In accordance with another aspect of the present invention, there are provided nucleic acid probes comprising nucleic acid molecules of sufficient length to
15 specifically hybridize to nucleic acid sequences of the present invention.

In accordance with another aspect of the present invention, there are provided methods of diagnosing diseases or a susceptibility to diseases related to mutations in nucleic acid sequences of the present invention and proteins encoded by such nucleic acid sequences.

20 In accordance with yet a further aspect of the present invention, there is provided a process for utilizing such polypeptides, or polynucleotides encoding such polypeptides, for *in vitro* purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors.

25 These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

Brief Description of the Figures

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

30 Figures 1A-1E show the full length nucleotide (SEQ ID NO:1) and the deduced amino acid (SEQ ID NO:2) sequence of VEGF-2. The polypeptide comprises approximately 419 amino acid residues of which approximately 23 represent the leader sequence. The standard one letter abbreviations for amino acids are used. Sequencing was performed using the Model 373 Automated DNA
35 Sequencer (Applied Biosystems, Inc.). Sequencing accuracy is predicted to be greater than 97%.

Figures 2A-2D show the nucleotide (SEQ ID NO:3) and the deduced amino acid (SEQ ID NO:4) sequence of a truncated, biologically active form of VEGF-2. The polypeptide comprises approximately 350 amino acid residues of which approximately the first 24 amino acids represent the leader sequence.

5 Figures 3A-3B are an illustration of the amino acid sequence homology between PDGFa (SEQ ID NO:5), PDGFb (SEQ ID NO:6), VEGF (SEQ ID NO:7), and VEGF-2 (SEQ ID NO:4). The boxed areas indicate the conserved sequences and the location of the eight conserved cysteine residues.

10 Figure 4 shows, in table-form, the percent homology between PDGFa, PDGFb, VEGF, and VEGF-2.

Figure 5 shows the presence of VEGF-2 mRNA in human breast tumor cell lines.

Figure 6 depicts the results of a Northern blot analysis of VEGF-2 in human adult tissues.

15 Figure 7 shows a photograph of an SDS-PAGE gel after *in vitro* transcription, translation and electrophoresis of the polypeptide of the present invention. Lane 1: ¹⁴C and rainbow M.W. marker; Lane 2: FGF control; Lane 3: VEGF-2 produced by M13-reverse and forward primers; Lane 4: VEGF-2 produced by M13 reverse and VEGF-F4 primers; Lane 5: VEGF-2 produced by M13 reverse and VEGF-F5
20 primers.

Figures 8A and 8B depict photographs of SDS-PAGE gels. VEGF-2 polypeptide was expressed in a baculovirus system consisting of Sf9 cells. Protein from the medium and cytoplasm of cells were analyzed by SDS-PAGE under non-reducing (Figure 8A) and reducing (Figure 8B) conditions.

25 Figure 9 depicts a photograph of an SDS-PAGE gel. The medium from Sf9 cells infected with a nucleic acid sequence of the present invention was precipitated. The resuspended precipitate was analyzed by SDS-PAGE and stained with Coomassie brilliant blue.

30 Figure 10 depicts a photograph of an SDS-PAGE gel. VEGF-2 was purified from the medium supernatant and analyzed by SDS-PAGE in the presence or absence of the reducing agent b-mercaptoethanol and stained by Coomassie brilliant blue.

Figure 11 depicts reverse phase HPLC analysis of purified VEGF-2 using a RP-300 column (0.21 x 3 cm, Applied Biosystems, Inc.). The column was equilibrated with 0.1% trifluoroacetic acid (Solvent A) and the proteins eluted with a
35 7.5 min gradient from 0 to 60% Solvent B, composed of acetonitrile containing

0.07% TFA. The protein elution was monitored by absorbance at 215 nm ("red" line) and 280 nm ("blue" line). The percentage of Solvent B is shown by the "green" line.

Figure 12 is a bar graph illustrating the effect of partially-purified VEGF-2 protein on the growth of vascular endothelial cells in comparison to basic fibroblast growth factor.

Figure 13 is a bar graph illustrating the effect of purified VEGF-2 protein on the growth of vascular endothelial cells.

Figure 14 depicts expression of VEGF-2 mRNA in human fetal and adult tissues.

Figure 15 depicts expression of VEGF-2 mRNA in human primary culture cells.

Figure 16 depicts transient expression of VEGF-2 protein in COS-7 cells.

Figure 17 depicts VEGF-2 stimulated proliferation of human umbilical vein endothelial cells (HUVEC).

Figure 18 depicts VEGF-2 stimulated proliferation of dermal microvascular endothelial cells.

Figure 19 depicts the stimulatory effect of VEGF-2 on proliferation of microvascular, umbilical cord, endometrial, and bovine aortic endothelial cells.

Figure 20 depicts inhibition of PDGF-induced vascular (human aortic) smooth muscle cell proliferation.

Figure 21 depicts stimulation of migration of HUVEC and bovine microvascular endothelial cells (BMEC) by VEGF-2.

Figure 22 depicts stimulation of nitric oxide release of HUVEC by VEGF-2 and VEGF-1.

Figure 23 depicts inhibition of cord formation of microvascular endothelial cells (CADMEC) by VEGF-2.

Figure 24 depicts stimulation of angiogenesis by VEGF, VEGF-2, and bFGF in the CAM assay.

Figure 25 depicts restoration of certain parameters in the ischemic limb by VEGF-2 protein (Figure 25, top panels) and naked expression plasmid (Figure 25, middle panels): BP ratio (Figure 25a); Blood Flow and Flow Reserve (Figure 25b); Angiographic Score (Figure 25c); Capillary density (Figure 25d).

Figures 26 A-G depicts ability of VEGF-2 to affect the diastolic blood pressure in spontaneously hypertensive rats (SHR). Figures 26a and b depict the dose-dependent decrease in diastolic blood pressure achieved with VEGF-2. (Figures 26c and d depict the decreased mean arterial pressure (MAP) observed with VEGF-2.

Panel E shows the effect of increasing doses of VEGF-2 on the mean arterial pressure (MAP) of SHR rats. Panel F shows the effect of VEGF-2 on the diastolic pressure of SHR rats. Panel G shows the effect of VEGF-2 on the diastolic blood pressure of SHR rats.

5 Figure 27 depicts inhibition of VEGF-2N= and VEGF-2-induced proliferation.

10 Figure 28 shows a schematic representation of the pHE4a expression vector (SEQ ID NO:16). The locations of the kanamycin resistance marker gene, the multiple cloning site linker region, the oriC sequence, and the *lacIq* coding sequence are indicated.

15 Figure 29 shows the nucleotide sequence of the regulatory elements of the pHE4a promoter (SEQ ID NO:17). The two *lac* operator sequences, the Shine-Delgarno sequence (S/D), and the terminal *HindIII* and *NdeI* restriction sites (italicized) are indicated.

Detailed Description of the Preferred Embodiments

20 In accordance with one aspect of the present invention, there are provided isolated nucleic acid molecules comprising a polynucleotide encoding a VEGF-2 polypeptide having the deduced amino acid sequence of Figure 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. The nucleotide sequence shown in SEQ ID NO:1 was obtained by sequencing a cDNA clone, which was deposited on May 12, 1995 at the American Type Tissue Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, and given ATCC Deposit No. 97149.

25 In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules comprising a polynucleotide encoding a truncated VEGF-2 polypeptide having the deduced amino acid sequence of Figure 2 (SEQ ID NO:4), which was determined by sequencing a cloned cDNA. The nucleotide sequence shown in SEQ ID NO:3 was obtained by sequencing a cDNA clone, which was deposited on March 4, 1994 at the American Type Tissue Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, and given ATCC Deposit Number 75698.

30 Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were

35

predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

A polynucleotide encoding a polypeptide of the present invention may be obtained from early stage human embryo (week 8 to 9) osteoclastomas, adult heart or several breast cancer cell lines. The polynucleotide of this invention was discovered in a cDNA library derived from early stage human embryo week 9. It is structurally related to the VEGF/PDGF family. It contains an open reading frame encoding a protein of about 419 amino acid residues of which approximately the first 23 amino acid residues are the putative leader sequence such that the mature protein comprises 396 amino acids, and which protein exhibits the highest amino acid sequence homology to human vascular endothelial growth factor (30% identity), followed by PDGFa (24%) and PDGFb (22%). (See Figure 4). It is particularly important that all eight cysteines are conserved within all four members of the family (see boxed areas of Figure 3). In addition, the signature for the PDGF/VEGF family, PXCXXXXRCXGCCN, (SEQ ID NO:8) is conserved in VEGF-2 (see Figure 3). The homology between VEGF-2, VEGF and the two PDGFs is at the protein sequence level. No nucleotide sequence homology can be detected, and therefore, it would be difficult to isolate the VEGF-2 through simple approaches such as low stringency hybridization.

The VEGF-2 polypeptide of the present invention is meant to include the full length polypeptide and polynucleotide sequence which encodes for any leader sequences and for active fragments of the full length polypeptide. Active fragments are meant to include any portions of the full length amino acid sequence which have less than the full 419 amino acids of the full length amino acid sequence as shown in

SEQ ID NO:2, but still contain the eight cysteine residues shown conserved in Figure 3 and that still have VEGF-2 activity.

There are at least two alternatively spliced VEGF-2 mRNA sequences present in normal tissues. The two bands in Figure 7, lane 5 indicate the presence of the
5 alternatively spliced mRNA encoding the VEGF-2 polypeptide of the present invention.

The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be
10 the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature polypeptide may be identical to the coding sequence shown in Figure 1 or Figure 2, or that of the deposited clones, or may be a different coding sequence which, as a result of the redundancy or degeneracy of the genetic code, encodes the same, mature polypeptide as the DNA of Figure 1, Figure 2, or the
15 deposited cDNAs.

The polynucleotide which encodes for the mature polypeptide of Figure 1 or Figure 2 or for the mature polypeptides encoded by the deposited cDNAs may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequences such as a leader or secretory
20 sequence or a proprotein sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequences) and non-coding sequences, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

Thus, the term "polynucleotide encoding a polypeptide" encompasses a
25 polynucleotide which includes only coding sequences for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequences.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs, and derivatives of the polypeptide having the deduced amino acid sequence of Figures 1 or 2, or the
30 polypeptide encoded by the cDNA of the deposited clones. The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

Thus, the present invention includes polynucleotides encoding the same mature polypeptide as shown in Figures 1 or 2 or the same mature polypeptide
35 encoded by the cDNA of the deposited clones as well as variants of such polynucleotides which variants encode for a fragment, derivative, or analog of the

polypeptides of Figures 1 or 2; or the polypeptide encoded by the cDNA of the deposited clones. Such nucleotide variants include deletion variants, substitution variants, and addition or insertion variants.

As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figures 1 or 2, or of the coding sequence of the deposited clones. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

The present invention also includes polynucleotides, wherein the coding sequence for the mature polypeptide may be fused in the same reading frame to a polynucleotide which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

Thus, for example, the polynucleotide of the present invention may encode for a mature protein, or for a protein having a prosequence or for a protein having both a prosequence and presequence (leader sequence).

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, *e.g.* COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., *et al.*, *Cell* 37:767 (1984)).

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95% identical, and more preferably at least 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding

the polypeptide having the amino acid sequence at positions from about 1 to about 396 in SEQ ID NO:2; (d) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.97149; (e) a nucleotide sequence encoding the mature VEGF-2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.97149; or (f) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), or (e).

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95% identical, and more preferably at least 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:4; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:4, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the polypeptide having the amino acid sequence at positions from about 1 to about 326 in SEQ ID NO:4; (d) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.75698; (e) a nucleotide sequence encoding the mature VEGF-2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.75698; or (f) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), or (e).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a VEGF-2 polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the VEGF-2 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence

shown in SEQ ID NOS:1 or 3, or to the nucleotides sequence of the deposited cDNA clone(s) can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

As described in detail below, the polypeptides of the present invention can be used to raise polyclonal and monoclonal antibodies, which are useful in diagnostic assays for detecting VEGF-2 protein expression as described below or as agonists and antagonists capable of enhancing or inhibiting VEGF-2 protein function. Further, such polypeptides can be used in the yeast two-hybrid system to "capture" VEGF-2 protein binding proteins which are also candidate agonist and antagonist according to the present invention. The yeast two hybrid system is described in Fields and Song, *Nature* 340:245-246 (1989).

In another aspect, the invention provides a peptide or polypeptide comprising an epitope-bearing portion of a polypeptide of the invention. As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, Sutcliffe, J. G., Shinnick, T. M., Green, N. and Learner, R. A. (1983) Antibodies that react with predetermined sites on proteins. *Science* 219:660-666. Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals. Peptides that are extremely hydrophobic and those of six or fewer residues generally are ineffective at inducing antibodies that bind to the mimicked protein; longer, soluble peptides, especially those containing proline residues, usually are effective. Sutcliffe *et al.*, *supra*, at 661. For instance, 18 of 20 peptides designed according to these guidelines,

containing 8-39 residues covering 75% of the sequence of the influenza virus hemagglutinin HA1 polypeptide chain, induced antibodies that reacted with the HA1 protein or intact virus; and 12/12 peptides from the MuLV polymerase and 18/18 from the rabies glycoprotein induced antibodies that precipitated the respective proteins.

Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind specifically to a polypeptide of the invention. Thus, a high proportion of hybridomas obtained by fusion of spleen cells from donors immunized with an antigen epitope-bearing peptide generally secrete antibody reactive with the native protein. Sutcliffe *et al.*, supra, at 663. The antibodies raised by antigenic epitope-bearing peptides or polypeptides are useful to detect the mimicked protein, and antibodies to different peptides may be used for tracking the fate of various regions of a protein precursor which undergoes post-translational processing. The peptides and anti-peptide antibodies may be used in a variety of qualitative or quantitative assays for the mimicked protein, for instance in competition assays since it has been shown that even short peptides (e.g., about 9 amino acids) can bind and displace the larger peptides in immunoprecipitation assays. See, for instance, Wilson *et al.*, *Cell* 37:767-778 (1984) at 777. The anti-peptide antibodies of the invention also are useful for purification of the mimicked protein, for instance, by adsorption chromatography using methods well known in the art.

Antigenic epitope-bearing peptides and polypeptides of the invention designed according to the above guidelines preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention. However, peptides or polypeptides comprising a larger portion of an amino acid sequence of a polypeptide of the invention, containing about 30, 40, 50, 60, 70, 80, 90, 100, or 150 amino acids, or any length up to and including the entire amino acid sequence of a polypeptide of the invention, also are considered epitope-bearing peptides or polypeptides of the invention and also are useful for inducing antibodies that react with the mimicked protein. Preferably, the amino acid sequence of the epitope-bearing peptide is selected to provide substantial solubility in aqueous solvents (i.e., the sequence includes relatively hydrophilic residues and highly hydrophobic sequences are preferably avoided); and sequences containing proline residues are particularly preferred.

Non-limiting examples of antigenic polypeptides or peptides that can be used to generate VEGF-2-specific antibodies include the following: a polypeptide comprising amino acid residues from about leu-37 to about glu-45 in SEQ ID NO:2, from about Tyr-58 to about Gly-66 in SEQ ID NO:2, from about Gln-73 to about Glu-81 in SEQ ID NO:2, from about Asp-100 to about Cys-108 in SEQ ID NO:2, from about Gly-140 to about Leu-148 in SEQ ID NO:2, from about Pro-168 to about Val-176 in SEQ ID NO:2, from about His-183 to about Lys-191 in SEQ ID NO:2, from about Ile-201 to about Thr-209 in SEQ ID NO:2, from about Ala-216 to about Tyr-224 in SEQ ID NO:2, from about Asp-244 to about His-254 in SEQ ID NO:2, from about Gly-258 to about Glu-266 in SEQ ID NO:2, from about Cys-272 to about Ser-280 in SEQ ID NO:2, from about Pro-283 to about Ser-291 in SEQ ID NO:2, from about Cys-296 to about Gln-304 in SEQ ID NO:2, from about Ala-307 to about Cys-316 in SEQ ID NO:2, from about Val-319 to about Cys-335 in SEQ ID NO:2, from about Cys-339 to about Leu-347 in SEQ ID NO:2, from about Cys-360 to about Glu-373 in SEQ ID NO:2, from about Tyr-378 to about Val-386 in SEQ ID NO:2, and from about Ser-388 to about Ser-396 in SEQ ID NO:2. These polypeptide fragments have been determined to bear antigenic epitopes of the VEGF-2 protein by the analysis of the Jameson-Wolf antigenic index.

The epitope-bearing peptides and polypeptides of the invention may be produced by any conventional means for making peptides or polypeptides including recombinant means using nucleic acid molecules of the invention. For instance, a short epitope-bearing amino acid sequence may be fused to a larger polypeptide that acts as a carrier during recombinant production and purification, as well as during immunization to produce anti-peptide antibodies. Epitope-bearing peptides also may be synthesized using known methods of chemical synthesis. For instance, Houghten has described a simple method for synthesis of large numbers of peptides, such as 10-20 mg of 248 different 13 residue peptides representing single amino acid variants of a segment of the HA1 polypeptide which were prepared and characterized (by ELISA-type binding studies) in less than four weeks. Houghten, R. A. (1985) General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 82:5131-5135. This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten *et al.* (1986). In this procedure the individual resins for the solid-phase synthesis of various peptides are contained in separate solvent-permeable packets, enabling the optimal use of the many identical repetitive steps involved in solid-phase methods. A completely

manual procedure allows 500-1000 or more syntheses to be conducted simultaneously. Houghten *et al.*, supra, at 5134.

Epitope-bearing peptides and polypeptides of the invention are used to induce antibodies according to methods well known in the art. See, for instance, Sutcliffe *et al.*, supra; Wilson *et al.*, supra; Chow, M. *et al.*, *Proc. Natl. Acad. Sci. USA* 82:910-914; and Bittle, F. J. *et al.*, *J. Gen. Virol.* 66:2347-2354 (1985). Generally, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling of the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine may be coupled to carrier using a linker such as m-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carrier using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier-coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 mg peptide or carrier protein and Freund's adjuvant. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

Immunogenic epitope-bearing peptides of the invention, i.e., those parts of a protein that elicit an antibody response when the whole protein is the immunogen, are identified according to methods known in the art. For instance, Geysen *et al.*, supra, discloses a procedure for rapid concurrent synthesis on solid supports of hundreds of peptides of sufficient purity to react in an enzyme-linked immunosorbent assay. Interaction of synthesized peptides with antibodies is then easily detected without removing them from the support. In this manner a peptide bearing an immunogenic epitope of a desired protein may be identified routinely by one of ordinary skill in the art. For instance, the immunologically important epitope in the coat protein of foot-and-mouth disease virus was located by Geysen *et al.* with a resolution of seven amino acids by synthesis of an overlapping set of all 208 possible hexapeptides covering the entire 213 amino acid sequence of the protein. Then, a complete replacement set of peptides in which all 20 amino acids were substituted in turn at every position within the epitope were synthesized, and the particular amino acids conferring specificity for the reaction with antibody were determined. Thus, peptide

analog of the epitope-bearing peptides of the invention can be made routinely by this method. U.S. Patent No. 4,708,781 to Geysen (1987) further describes this method of identifying a peptide bearing an immunogenic epitope of a desired protein.

Further still, U.S. Patent No. 5,194,392 to Geysen (1990) describes a
5 general method of detecting or determining the sequence of monomers (amino acids or other compounds) which is a topological equivalent of the epitope (i.e., a Amimotope) which is complementary to a particular paratope (antigen binding site) of an antibody of interest. More generally, U.S. Patent No. 4,433,092 to Geysen (1989) describes a
10 method of detecting or determining a sequence of monomers which is a topographical equivalent of a ligand which is complementary to the ligand binding site of a particular receptor of interest. Similarly, U.S. Patent No. 5,480,971 to Houghten, R. A. *et al.* (1996) on Peralkylated Oligopeptide Mixtures discloses linear C₁-C₇-alkyl
15 peralkylated oligopeptides and sets and libraries of such peptides, as well as methods for using such oligopeptide sets and libraries for determining the sequence of a peralkylated oligopeptide that preferentially binds to an acceptor molecule of interest. Thus, non-peptide analogs of the epitope-bearing peptides of the invention also can be made routinely by these methods.

As one of skill in the art will appreciate, VEGF-2 polypeptides of the present invention and the epitope-bearing fragments thereof described above can be combined
20 with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EPA 394,827;
25 Traunecker *et al.*, *Nature* 331:84- 86 (1988)).

In accordance with the present invention, novel variants of VEGF-2 are also described. These can be produced by deleting or substituting one or more amino acids of VEGF-2. Natural mutations are called allelic variations. Allelic variations can be
30 silent (no change in the encoded polypeptide) or may have altered amino acid sequence.

In order to attempt to improve or alter the characteristics of native VEGF-2, protein engineering may be employed. Recombinant DNA technology known to those skilled in the art can be used to create novel polypeptides. Muteins and deletions can show, e.g., enhanced activity or increased stability. In addition, they could be purified
35 in higher yield and show better solubility at least under certain purification and storage conditions. Set forth below are examples of mutations that can be constructed.

Amino terminal and carboxy terminal deletions

Furthermore, VEGF-2 appears to be proteolytically cleaved upon expression resulting in polypeptide fragments of the following sizes when run on a SDS-PAGE gel (sizes are approximate) (See, Figures 6-8, for example): 80, 59, 45, 43, 41, 40, 39, 38, 37, 36, 31, 29, 21, and 15 kDa. These polypeptide fragments are the result of proteolytic cleavage at both the N-terminal and C-terminal portions of the protein. These proteolytically generated fragments appears to have activity, particularly the 21 kDa fragment.

In addition, protein engineering may be employed in order to improve or alter one or more characteristics of native VEGF-2. The deletion of carboxyterminal amino acids can enhance the activity of proteins. One example is interferon gamma that shows up to ten times higher activity by deleting ten amino acid residues from the carboxy terminus of the protein (Döbeli *et al.*, *J. of Biotechnology* 7:199-216 (1988)). Thus, one aspect of the invention is to provide polypeptide analogs of VEGF-2 and nucleotide sequences encoding such analogs that exhibit enhanced stability (e.g., when exposed to typical pH, thermal conditions or other storage conditions) relative to the native VEGF-2 polypeptide.

Particularly preferred VEGF-2 polypeptides are shown below (numbering starts with the first amino acid in the protein (Met) (Figure 1 (SEQ ID NO:18)): Ala (residue 24) to Ser (residue 419); Pro (25) to Ser (419); Ala (26) to Ser (419); Ala (27) to Ser (419); Ala (28) to Ser (419); Ala (29) to Ser (419); Ala (30) to Ser (419); Phe (31) to Ser (419); Glu (32) to Ser (419); Ser (33) to Ser (419); Gly (34) to Ser (419); Leu (35) to Ser (419); Asp (36) to Ser (419); Leu (37) to (Ser (419); Ser (38) to Ser (419); Asp (39) to Ser (419); Ala (40) to Ser (419); Glu (41) to Ser (419); Pro (42) to Ser (419); Asp (43) to Ser (419); Ala (44) to Ser (419); Gly (45) to Ser (419); Glu (46) to Ser (419); Ala (47) to Ser (419); Thr (48) to Ser (419); Ala (49) to Ser (419); Tyr (50) to Ser (419); Ser (52) to Ser (419); Asp (54) to Ser (419); Val (62) to Ser (419); Val (65) to Ser (419); Met(1), Glu (23), or Ala (24) to Met (418); Met (1), Glu (23), or Ala (24) to Gln (417); Met (1), Glu (23), or Ala (24) to Pro (416); Met(1), Glu (23), or Ala (24) to Arg (415); Met(1), Glu (23), or Ala (24) to Gln (414); Met(1), Glu (23), or Ala (24) to Trp (413); Met(1), Glu (23), or Ala (24) to Tyr (412); Met(1), Glu (23), or Ala (24) to Ser (411); Met(1), Glu (23), or Ala (24) to Pro (410); Met (1), Glu (23), or Ala(24) to Val (409); Met (1), Glu (23), or Ala (24) to Cys (408); Met(1), Glu (23), or Ala (24) to Arg (407); Met(1), Glu (23), or Ala (24) to Cys (406); Met (1), Glu (23), or Ala (24) to Val (405); Met(1), Glu (23), or Ala (24) to Glu (404); Met(1), Glu (23), or Ala (24) to Glu (403); Met(1), Glu

(23), or Ala (24) to Ser (402); Met(1), Glu (23), or Ala (24) to Gly (398); Met(1), Glu (23), or Ala (24) to Pro (397); Met(1), Glu (23), or Ala (24) to Lys (393); Met(1), Glu (23), or Ala (24) to Met(263); Met(1), Glu (23), or Ala (24) to Asp(311); Met(1), Glu (23), or Ala (24) to Pro (367); Met(1) to Ser (419); Met(1) to Ser(228); Glu(47) to Ser(419); Ala(111) to Lys(214); Ala(112) to Lys(214); His(113) to Lys(214); Tyr(114) to Lys(214); Asn(115) to Lys(214); Thr(116) to Lys(214); Thr(103) to Leu(215); Glu(104) to Leu(215); Glu(105) to Leu(215); Thr(106) to Leu(215); Ile(107) to Leu(215); Lys(108) to Leu(215); Phe(109) to Leu(215); Ala(110) to Leu(215); Ala(111) to Leu(215); Ala(112) to Leu(215); His(113) to Leu(215); Tyr(114) to Leu(215); Asn(115) to Leu(215); Thr(116) to Leu(215); Thr(103) to Ser(228); Glu(104) to Ser(228); Glu(105) to Ser(228); Thr(106) to Ser(228); Ile(107) to Ser(228); Lys(108) to Ser(228); Phe(109) to Ser(228); Ala(110) to Ser(228); Ala(111) to Ser(228); Ala(112) to Ser(228); His(113) to Ser(228); Tyr(114) to Ser(228); Asn(115) to Ser(228); Thr(116) to Ser(228); Thr(103) to Leu(229); Glu(104) to Leu(229); Thr(103) to Arg(227); Glu(104) to Arg(227); Glu(105) to Arg (227); Thr(106) to Arg (227); Ile(107) to Arg (227); Lys(108) to Arg (227); Phe(109) to Arg (227); Ala(110) to Arg (227); Ala(111) to Arg (227); Ala(112) to Arg (227); His(113) to Arg (227); Tyr(114) to Arg (227); Asn(115) to Arg (227); Thr(116) to Arg (227); Thr(103) to Ser(213); Glu(104) to Ser(213); Glu(105) to Ser(213); Thr(106) to Ser(213); Ile(107) to Ser(213); Lys(108) to Ser(213); Phe(109) to Ser(213); Ala(110) to Ser(213); Ala(111) to Ser(213); Ala(112) to Ser(213); His(113) to Ser(213); Tyr(114) to Ser(213); Asn(115) to Ser(213); Thr(116) to Ser(213); Thr(103) to Lys(214); Glu(104) to Lys(214); Glu(105) to Lys(214); Thr(106) to Lys(214); Ile(107) to Lys(214); Lys(108) to Lys(214); Phe(109) to Lys(214); Ala(110) to Lys(214); Glu(105) to Leu(229); Thr(106) to Leu(229); Ile(107) to Leu(229); Lys(108) to Leu(229); Phe(109) to Leu(229); Ala(110) to Leu(229); Ala(111) to Leu(229); Ala(112) to Leu(229); His(113) to Leu(229); Tyr(114) to Leu(229); Asn(115) to Leu(229); Thr(116) to Leu(229).

Preferred embodiments include the following deletion mutants: Thr(103) -- Arg(227); Glu(104) -- Arg(227); Ala(112) -- Arg (227); Thr(103) -- Ser(213); Glu(104) -- Ser(213); Thr(103) -- Leu(215); Glu(47) -- Ser(419); Met(1), Glu (23), or Ala (24) -- Met(263); Met(1), Glu (23), or Ala (24) -- Asp(311); Met(1), Glu (23), or Ala (24) -- Pro (367); Met(1) -- Ser(419); and Met(1) -- Ser(228) of (Figure 1 (SEQ ID NO:18)).

Also included by the present invention are deletion mutants having amino acids deleted from both the NB terminus and the C-terminus. Such mutants include all combinations of the N-terminal deletion mutants and C-terminal deletion mutants described above. Those combinations can be made using recombinant techniques known to those skilled in the art.

Particularly, N-terminal deletions of the VEGF-2 polypeptide can be described by the general formula m-396, where m is an integer from -23 to 388, where m corresponds to the position of the amino acid residue identified in SEQ ID NO:2. Preferably, N-terminal deletions retain the conserved boxed area of Figure 3 (PXC~~V~~XXXRCXGCCN)(SEQ ID NO: 8), and include polypeptides comprising the amino acid sequence of residues: N-terminal deletions of the polypeptide of the invention shown as SEQ ID NO:1 include polypeptides comprising the amino acid sequence of residues: : E-1 to S-396; A-2 to S-396; P-3 to S-396; A-4 to S-396; A-5 to S-396; A-6 to S-396; A-7 to S-396; A-8 to S-396; F-9 to S-396; E-10 to S-396; S-11 to S-396; G-12 to S-396; L-13 to S-396; D-14 to S-396; L-15 to S-396; S-16 to S-396; D-17 to S-396; A-18 to S-396; E-19 to S-396; P-20 to S-396; D-21 to S-396; A-22 to S-396; G-23 to S-396; E-24 to S-396; A-25 to S-396; T-26 to S-396; A-27 to S-396; Y-28 to S-396; A-29 to S-396; S-30 to S-396; K-31 to S-396; D-32 to S-396; L-33 to S-396; E-34 to S-396; E-35 to S-396; Q-36 to S-396; L-37 to S-396; R-38 to S-396; S-39 to S-396; V-40 to S-396; S-41 to S-396; S-42 to S-396; V-43 to S-396; D-44 to S-396; E-45 to S-396; L-46 to S-396; M-47 to S-396; T-48 to S-396; V-49 to S-396; L-50 to S-396; Y-51 to S-396; P-52 to S-396; E-53 to S-396; Y-54 to S-396; W-55 to S-396; K-56 to S-396; M-57 to S-396; Y-58 to S-396; K-59 to S-396; C-60 to S-396; Q-61 to S-396; L-62 to S-396; R-63 to S-396; K-64 to S-396; G-65 to S-396; G-66 to S-396; W-67 to S-396; Q-68 to S-396; H-69 to S-396; N-70 to S-396; R-71 to S-396; E-72 to S-396; Q-73 to S-396; A-74 to S-396; N-75 to S-396; L-76 to S-396; N-77 to S-396; S-78 to S-396; R-79 to S-396; T-80 to S-396; E-81 to S-396; E-82 to S-396; T-83 to S-396; I-84 to S-396; K-85 to S-396; F-86 to S-396; A-87 to S-396; A-88 to S-396; A-89 to S-396; H-90 to S-396; Y-91 to S-396; N-92 to S-396; T-93 to S-396; E-94 to S-396; I-95 to S-396; L-96 to S-396; K-97 to S-396; S-98 to S-396; I-99 to S-396; D-100 to S-396; N-101 to S-396; E-102 to S-396; W-103 to S-396; R-104 to S-396; K-105 to S-396; T-106 to S-396; Q-107 to S-396; C-108 to S-396; M-109 to S-396; P-110 to S-396; R-111 to S-396; E-112 to S-396; V-113 to S-396; C-114 to S-396; I-115 to S-396; D-116 to S-396; V-117 to S-396; G-118 to S-396; K-119 to S-396; E-120 to S-396; F-121 to S-396; G-122 to S-396; V-123 to S-396; A-124 to S-396; T-125 to S-396; N-126 to S-396; T-127 to S-396; F-128 to S-

396; F-129 to S-396; K-130 to S-396; P-131 to S-396; P-132 to S-396; C-133 to S-396; V-134 to S-396; S-135 to S-396; V-136 to S-396; Y-137 to S-396; R-138 to S-396; C-139 to S-396; G-140 to S-396; G-141 to S-396; C-142 to S-396; C-143 to S-396; N-144 to S-396; S-145 to S-396; E-146 to S-396; G-147 to S-396; L-148 to S-396; Q-149 to S-396; C-150 to S-396; M-151 to S-396; N-152 to S-396; T-153 to S-396; S-154 to S-396; T-155 to S-396; S-156 to S-396; Y-157 to S-396; L-158 to S-396; S-159 to S-396; K-160 to S-396; T-161 to S-396; L-162 to S-396; F-163 to S-396; E-164 to S-396; I-165 to S-396; T-166 to S-396; V-167 to S-396; P-168 to S-396; L-169 to S-396; S-170 to S-396; Q-171 to S-396; G-172 to S-396; P-173 to S-396; K-174 to S-396; P-175 to S-396; V-176 to S-396; T-177 to S-396; I-178 to S-396; S-179 to S-396; F-180 to S-396; A-181 to S-396; N-182 to S-396; H-183 to S-396; T-184 to S-396; S-185 to S-396; C-186 to S-396; R-187 to S-396; C-188 to S-396; M-189 to S-396; S-190 to S-396; K-191 to S-396; L-192 to S-396; D-193 to S-396; V-194 to S-396; Y-195 to S-396; R-196 to S-396; Q-197 to S-396; V-198 to S-396; H-199 to S-396; S-200 to S-396; I-201 to S-396; I-202 to S-396; R-203 to S-396; R-204 to S-396; S-205 to S-396; L-206 to S-396; P-207 to S-396; A-208 to S-396; T-209 to S-396; L-210 to S-396; P-211 to S-396; Q-212 to S-396; C-213 to S-396; Q-214 to S-396; A-215 to S-396; A-216 to S-396; N-217 to S-396; K-218 to S-396; T-219 to S-396; C-220 to S-396; P-221 to S-396; T-222 to S-396; N-223 to S-396; Y-224 to S-396; M-225 to S-396; W-226 to S-396; N-227 to S-396; N-228 to S-396; H-229 to S-396; I-230 to S-396; C-231 to S-396; R-232 to S-396; C-233 to S-396; L-234 to S-396; A-235 to S-396; Q-236 to S-396; E-237 to S-396; D-238 to S-396; F-239 to S-396; M-240 to S-396; F-241 to S-396; S-242 to S-396; S-243 to S-396; D-244 to S-396; A-245 to S-396; G-246 to S-396; D-247 to S-396; D-248 to S-396; S-249 to S-396; T-250 to S-396; D-251 to S-396; G-252 to S-396; F-253 to S-396; H-254 to S-396; D-255 to S-396; I-256 to S-396; C-257 to S-396; G-258 to S-396; P-259 to S-396; N-260 to S-396; K-261 to S-396; E-262 to S-396; L-263 to S-396; D-264 to S-396; E-265 to S-396; E-266 to S-396; T-267 to S-396; C-268 to S-396; Q-269 to S-396; C-270 to S-396; V-271 to S-396; C-272 to S-396; R-273 to S-396; A-274 to S-396; G-275 to S-396; L-276 to S-396; R-277 to S-396; P-278 to S-396; A-279 to S-396; S-280 to S-396; C-281 to S-396; G-282 to S-396; P-283 to S-396; H-284 to S-396; K-285 to S-396; E-286 to S-396; L-287 to S-396; D-288 to S-396; R-289 to S-396; N-290 to S-396; S-291 to S-396; C-292 to S-396; Q-293 to S-396; C-294 to S-396; V-295 to S-396; C-296 to S-396; K-297 to S-396; N-298 to S-396; K-299 to S-396; L-300 to S-396; F-301 to S-396; P-302 to S-396; S-303 to S-396; Q-304 to S-396; C-305 to S-396; G-306 to S-396; A-307 to S-396; N-308 to S-

396; R-309 to S-396; E-310 to S-396; F-311 to S-396; D-312 to S-396; E-313 to S-396; N-314 to S-396; T-315 to S-396; C-316 to S-396; Q-317 to S-396; C-318 to S-396; V-319 to S-396; C-320 to S-396; K-321 to S-396; R-322 to S-396; T-323 to S-396; C-324 to S-396; P-325 to S-396; R-326 to S-396; N-327 to S-396; Q-328 to S-396; P-329 to S-396; L-330 to S-396; N-331 to S-396; P-332 to S-396; G-333 to S-396; K-334 to S-396; C-335 to S-396; A-336 to S-396; C-337 to S-396; E-338 to S-396; C-339 to S-396; T-340 to S-396; E-341 to S-396; S-342 to S-396; P-343 to S-396; Q-344 to S-396; K-345 to S-396; C-346 to S-396; L-347 to S-396; L-348 to S-396; K-349 to S-396; G-350 to S-396; K-351 to S-396; K-352 to S-396; F-353 to S-396; H-354 to S-396; H-355 to S-396; Q-356 to S-396; T-357 to S-396; C-358 to S-396; S-359 to S-396; C-360 to S-396; Y-361 to S-396; R-362 to S-396; R-363 to S-396; P-364 to S-396; C-365 to S-396; T-366 to S-396; N-367 to S-396; R-368 to S-396; Q-369 to S-396; K-370 to S-396; A-371 to S-396; C-372 to S-396; E-373 to S-396; P-374 to S-396; G-375 to S-396; F-376 to S-396; S-377 to S-396; Y-378 to S-396; S-379 to S-396; E-380 to S-396; E-381 to S-396; V-382 to S-396; C-383 to S-396; R-384 to S-396; C-385 to S-396; V-386 to S-396; P-387 to S-396; S-388 to S-396; Y-389 to S-396; W-390 to S-396; Q-391 to S-396 of SEQ ID NO:2. One preferred embodiment comprises amino acids S-205 to S-396 of SEQ ID NO:2. Also preferred are polynucleotides encoding these polypeptides.

Moreover, C-terminal deletions of the VEGF-2 polypeptide can also be described by the general formula -23-n, where n is an integer from -15 to 395 where n corresponds to the position of amino acid residue identified in SEQ ID NO:2. Preferably, C-terminal deletions retain the conserved boxed area of Figure 3 (PXCXXXXRCXGCCN)(SEQ ID NO: 8), and include polypeptides comprising the amino acid sequence of residues: Likewise, C-terminal deletions of the polypeptide of the invention shown as SEQ ID NO:2 include polypeptides comprising the amino acid sequence of residues: E-1 to M-395; E-1 to Q-394; E-1 to P-393; E-1 to R-392; E-1 to Q-391; E-1 to W-390; E-1 to Y-389; E-1 to S-388; E-1 to P-387; E-1 to V-386; E-1 to C-385; E-1 to R-384; E-1 to C-383; E-1 to V-382; E-1 to E-381; E-1 to E-380; E-1 to S-379; E-1 to Y-378; E-1 to S-377; E-1 to F-376; E-1 to G-375; E-1 to P-374; E-1 to E-373; E-1 to C-372; E-1 to A-371; E-1 to K-370; E-1 to Q-369; E-1 to R-368; E-1 to N-367; E-1 to T-366; E-1 to C-365; E-1 to P-364; E-1 to R-363; E-1 to R-362; E-1 to Y-361; E-1 to C-360; E-1 to S-359; E-1 to C-358; E-1 to T-357; E-1 to Q-356; E-1 to H-355; E-1 to H-354; E-1 to F-353; E-1 to K-352; E-1 to K-351; E-1 to G-350; E-1 to K-349; E-1 to L-348; E-1 to L-347; E-1 to C-346; E-1 to K-345; E-1 to Q-344; E-1 to P-343; E-1 to S-342; E-1 to E-341; E-1 to T-340; E-1 to C-339; E-1 to E-338; E-1

to C-337; E-1 to A-336; E-1 to C-335; E-1 to K-334; E-1 to G-333; E-1 to P-332; E-1
to N-331; E-1 to L-330; E-1 to P-329; E-1 to Q-328; E-1 to N-327; E-1 to R-326; E-1
to P-325; E-1 to C-324; E-1 to T-323; E-1 to R-322; E-1 to K-321; E-1 to C-320; E-1
to V-319; E-1 to C-318; E-1 to Q-317; E-1 to C-316; E-1 to T-315; E-1 to N-314; E-1
5 to E-313; E-1 to D-312; E-1 to F-311; E-1 to E-310; E-1 to R-309; E-1 to N-308; E-1
to A-307; E-1 to G-306; E-1 to C-305; E-1 to Q-304; E-1 to S-303; E-1 to P-302; E-1
to F-301; E-1 to L-300; E-1 to K-299; E-1 to N-298; E-1 to K-297; E-1 to C-296; E-1
to V-295; E-1 to C-294; E-1 to Q-293; E-1 to C-292; E-1 to S-291; E-1 to N-290; E-1
to R-289; E-1 to D-288; E-1 to L-287; E-1 to E-286; E-1 to K-285; E-1 to H-284; E-1
10 to P-283; E-1 to G-282; E-1 to C-281; E-1 to S-280; E-1 to A-279; E-1 to P-278; E-1
to R-277; E-1 to L-276; E-1 to G-275; E-1 to A-274; E-1 to R-273; E-1 to C-272; E-1
to V-271; E-1 to C-270; E-1 to Q-269; E-1 to C-268; E-1 to T-267; E-1 to E-266; E-1
to E-265; E-1 to D-264; E-1 to L-263; E-1 to E-262; E-1 to K-261; E-1 to N-260; E-1
to P-259; E-1 to G-258; E-1 to C-257; E-1 to I-256; E-1 to D-255; E-1 to H-254; E-1
15 to F-253; E-1 to G-252; E-1 to D-251; E-1 to T-250; E-1 to S-249; E-1 to D-248; E-1
to D-247; E-1 to G-246; E-1 to A-245; E-1 to D-244; E-1 to S-243; E-1 to S-242; E-1
to F-241; E-1 to M-240; E-1 to F-239; E-1 to D-238; E-1 to E-237; E-1 to Q-236; E-1
to A-235; E-1 to L-234; E-1 to C-233; E-1 to R-232; E-1 to C-231; E-1 to I-230; E-1
to H-229; E-1 to N-228; E-1 to N-227; E-1 to W-226; E-1 to M-225; E-1 to Y-224; E-1
20 to N-223; E-1 to T-222; E-1 to P-221; E-1 to C-220; E-1 to T-219; E-1 to K-218; E-1
to N-217; E-1 to A-216; E-1 to A-215; E-1 to Q-214; E-1 to C-213; E-1 to Q-212;
E-1 to P-211; E-1 to L-210; E-1 to T-209; E-1 to A-208; E-1 to P-207; E-1 to L-206;
E-1 to S-205; E-1 to R-204; E-1 to R-203; E-1 to I-202; E-1 to I-201; E-1 to S-200;
E-1 to H-199; E-1 to V-198; E-1 to Q-197; E-1 to R-196; E-1 to Y-195; E-1 to V-194;
25 E-1 to D-193; E-1 to L-192; E-1 to K-191; E-1 to S-190; E-1 to M-189; E-1 to C-188;
E-1 to R-187; E-1 to C-186; E-1 to S-185; E-1 to T-184; E-1 to H-183; E-1 to N-182;
E-1 to A-181; E-1 to F-180; E-1 to S-179; E-1 to I-178; E-1 to T-177; E-1 to V-176;
E-1 to P-175; E-1 to K-174; E-1 to P-173; E-1 to G-172; E-1 to Q-171; E-1 to S-170;
E-1 to L-169; E-1 to P-168; E-1 to V-167; E-1 to T-166; E-1 to I-165; E-1 to E-164;
30 E-1 to F-163; E-1 to L-162; E-1 to T-161; E-1 to K-160; E-1 to S-159; E-1 to L-158;
E-1 to Y-157; E-1 to S-156; E-1 to T-155; E-1 to S-154; E-1 to T-153; E-1 to N-152;
E-1 to M-151; E-1 to C-150; E-1 to Q-149; E-1 to L-148; E-1 to G-147; E-1 to E-146;
E-1 to S-145; E-1 to N-144; E-1 to C-143; E-1 to C-142; E-1 to G-141; E-1 to G-140;
E-1 to C-139; E-1 to R-138; E-1 to Y-137; E-1 to V-136; E-1 to S-135; E-1 to V-134;
35 E-1 to C-133; E-1 to P-132; E-1 to P-131; E-1 to K-130; E-1 to F-129; E-1 to F-128;
E-1 to T-127; E-1 to N-126; E-1 to T-125; E-1 to A-124; E-1 to V-123; E-1 to G-122;

E-1 to F-121; E-1 to E-120; E-1 to K-119; E-1 to G-118; E-1 to V-117; E-1 to D-116; E-1 to I-115; E-1 to C-114; E-1 to V-113; E-1 to E-112; E-1 to R-111; E-1 to P-110; E-1 to M-109; E-1 to C-108; E-1 to Q-107; E-1 to T-106; E-1 to K-105; E-1 to R-104; E-1 to W-103; E-1 to E-102; E-1 to N-101; E-1 to D-100; E-1 to I-99; E-1 to S-98; E-1 to K-97; E-1 to L-96; E-1 to I-95; E-1 to E-94; E-1 to T-93; E-1 to N-92; E-1 to Y-91; E-1 to H-90; E-1 to A-89; E-1 to A-88; E-1 to A-87; E-1 to F-86; E-1 to K-85; E-1 to I-84; E-1 to T-83; E-1 to E-82; E-1 to E-81; E-1 to T-80; E-1 to R-79; E-1 to S-78; E-1 to N-77; E-1 to L-76; E-1 to N-75; E-1 to A-74; E-1 to Q-73; E-1 to E-72; E-1 to R-71; E-1 to N-70; E-1 to H-69; E-1 to Q-68; E-1 to W-67; E-1 to G-66; E-1 to G-65; E-1 to K-64; E-1 to R-63; E-1 to L-62; E-1 to Q-61; E-1 to C-60; E-1 to K-59; E-1 to Y-58; E-1 to M-57; E-1 to K-56; E-1 to W-55; E-1 to Y-54; E-1 to E-53; E-1 to P-52; E-1 to Y-51; E-1 to L-50; E-1 to V-49; E-1 to T-48; E-1 to M-47; E-1 to L-46; E-1 to E-45; E-1 to D-44; E-1 to V-43; E-1 to S-42; E-1 to S-41; E-1 to V-40; E-1 to S-39; E-1 to R-38; E-1 to L-37; E-1 to Q-36; E-1 to E-35; E-1 to E-34; E-1 to L-33; E-1 to D-32; E-1 to K-31; E-1 to S-30; E-1 to A-29; E-1 to Y-28; E-1 to A-27; E-1 to T-26; E-1 to A-25; E-1 to E-24; E-1 to G-23; E-1 to A-22; E-1 to D-21; E-1 to P-20; E-1 to E-19; E-1 to A-18; E-1 to D-17; E-1 to S-16; E-1 to L-15; E-1 to D-14; E-1 to L-13; E-1 to G-12; E-1 to S-11; E-1 to E-10; E-1 to F-9; E-1 to A-8; E-1 to A-7 of SEQ ID NO:2. Also preferred are polynucleotides encoding these polypeptides.

Moreover, the invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of SEQ ID NO:2, where n and m are integers as described above.

Likewise, also preferred are C-terminal deletions of the VEGF-2 polypeptide of the invention shown as SEQ ID NO:2 which include polypeptides comprising the amino acid sequence of residues: F-9 to M-395; F-9 to Q-394; F-9 to P-393; F-9 to R-392; F-9 to Q-391; F-9 to W-390; F-9 to Y-389; F-9 to S-388; F-9 to P-387; F-9 to V-386; F-9 to C-385; F-9 to R-384; F-9 to C-383; F-9 to V-382; F-9 to E-381; F-9 to E-380; F-9 to S-379; F-9 to Y-378; F-9 to S-377; F-9 to F-376; F-9 to G-375; F-9 to P-374; F-9 to E-373; F-9 to C-372; F-9 to A-371; F-9 to K-370; F-9 to Q-369; F-9 to R-368; F-9 to N-367; F-9 to T-366; F-9 to C-365; F-9 to P-364; F-9 to R-363; F-9 to R-362; F-9 to Y-361; F-9 to C-360; F-9 to S-359; F-9 to C-358; F-9 to T-357; F-9 to Q-356; F-9 to H-355; F-9 to H-354; F-9 to F-353; F-9 to K-352; F-9 to K-351; F-9 to G-350; F-9 to K-349; F-9 to L-348; F-9 to L-347; F-9 to C-346; F-9 to K-345; F-9 to Q-344; F-9 to P-343; F-9 to S-342; F-9 to E-341; F-9 to T-340; F-9 to C-339; F-9 to E-338; F-9 to C-337; F-9 to A-336; F-9 to C-335; F-9 to K-334; F-9 to G-333; F-9 to

P-332; F-9 to N-331; F-9 to L-330; F-9 to P-329; F-9 to Q-328; F-9 to N-327; F-9 to R-326; F-9 to P-325; F-9 to C-324; F-9 to T-323; F-9 to R-322; F-9 to K-321; F-9 to C-320; F-9 to V-319; F-9 to C-318; F-9 to Q-317; F-9 to C-316; F-9 to T-315; F-9 to N-314; F-9 to E-313; F-9 to D-312; F-9 to F-311; F-9 to E-310; F-9 to R-309; F-9 to
5 N-308; F-9 to A-307; F-9 to G-306; F-9 to C-305; F-9 to Q-304; F-9 to S-303; F-9 to P-302; F-9 to F-301; F-9 to L-300; F-9 to K-299; F-9 to N-298; F-9 to K-297; F-9 to C-296; F-9 to V-295; F-9 to C-294; F-9 to Q-293; F-9 to C-292; F-9 to S-291; F-9 to N-290; F-9 to R-289; F-9 to D-288; F-9 to L-287; F-9 to E-286; F-9 to K-285; F-9 to H-284; F-9 to P-283; F-9 to G-282; F-9 to C-281; F-9 to S-280; F-9 to A-279; F-9 to
10 P-278; F-9 to R-277; F-9 to L-276; F-9 to G-275; F-9 to A-274; F-9 to R-273; F-9 to C-272; F-9 to V-271; F-9 to C-270; F-9 to Q-269; F-9 to C-268; F-9 to T-267; F-9 to E-266; F-9 to E-265; F-9 to D-264; F-9 to L-263; F-9 to E-262; F-9 to K-261; F-9 to N-260; F-9 to P-259; F-9 to G-258; F-9 to C-257; F-9 to I-256; F-9 to D-255; F-9 to H-254; F-9 to F-253; F-9 to G-252; F-9 to D-251; F-9 to T-250; F-9 to S-249; F-9 to
15 D-248; F-9 to D-247; F-9 to G-246; F-9 to A-245; F-9 to D-244; F-9 to S-243; F-9 to S-242; F-9 to F-241; F-9 to M-240; F-9 to F-239; F-9 to D-238; F-9 to E-237; F-9 to Q-236; F-9 to A-235; F-9 to L-234; F-9 to C-233; F-9 to R-232; F-9 to C-231; F-9 to I-230; F-9 to H-229; F-9 to N-228; F-9 to N-227; F-9 to W-226; F-9 to M-225; F-9 to Y-224; F-9 to N-223; F-9 to T-222; F-9 to P-221; F-9 to C-220; F-9 to T-219; F-9
20 to K-218; F-9 to N-217; F-9 to A-216; F-9 to A-215; F-9 to Q-214; F-9 to C-213; F-9 to Q-212; F-9 to P-211; F-9 to L-210; F-9 to T-209; F-9 to A-208; F-9 to P-207; F-9 to L-206; F-9 to S-205; F-9 to R-204; F-9 to R-203; F-9 to I-202; F-9 to I-201; F-9 to S-200; F-9 to H-199; F-9 to V-198; F-9 to Q-197; F-9 to R-196; F-9 to Y-195; F-9 to V-194; F-9 to D-193; F-9 to L-192; F-9 to K-191; F-9 to S-190; F-9 to M-189; F-9 to
25 C-188; F-9 to R-187; F-9 to C-186; F-9 to S-185; F-9 to T-184; F-9 to H-183; F-9 to N-182; F-9 to A-181; F-9 to F-180; F-9 to S-179; F-9 to I-178; F-9 to T-177; F-9 to V-176; F-9 to P-175; F-9 to K-174; F-9 to P-173; F-9 to G-172; F-9 to Q-171; F-9 to S-170; F-9 to L-169; F-9 to P-168; F-9 to V-167; F-9 to T-166; F-9 to I-165; F-9 to E-164; F-9 to F-163; F-9 to L-162; F-9 to T-161; F-9 to K-160; F-9 to S-159; F-9 to
30 L-158; F-9 to Y-157; F-9 to S-156; F-9 to T-155; F-9 to S-154; F-9 to T-153; F-9 to N-152; F-9 to M-151; F-9 to C-150; F-9 to Q-149; F-9 to L-148; F-9 to G-147; F-9 to E-146; F-9 to S-145; F-9 to N-144; F-9 to C-143; F-9 to C-142; F-9 to G-141; F-9 to G-140; F-9 to C-139; F-9 to R-138; F-9 to Y-137; F-9 to V-136; F-9 to S-135; F-9 to V-134; F-9 to C-133; F-9 to P-132; F-9 to P-131; F-9 to K-130; F-9 to F-129; F-9 to
35 F-128; F-9 to T-127; F-9 to N-126; F-9 to T-125; F-9 to A-124; F-9 to V-123; F-9 to G-122; F-9 to F-121; F-9 to E-120; F-9 to K-119; F-9 to G-118; F-9 to V-117; F-9 to

D-116; F-9 to I-115; F-9 to C-114; F-9 to V-113; F-9 to E-112; F-9 to R-111; F-9 to P-110; F-9 to M-109; F-9 to C-108; F-9 to Q-107; F-9 to T-106; F-9 to K-105; F-9 to R-104; F-9 to W-103; F-9 to E-102; F-9 to N-101; F-9 to D-100; F-9 to I-99; F-9 to S-98; F-9 to K-97; F-9 to L-96; F-9 to I-95; F-9 to E-94; F-9 to T-93; F-9 to N-92; F-9 to Y-91; F-9 to H-90; F-9 to A-89; F-9 to A-88; F-9 to A-87; F-9 to F-86; F-9 to K-85; F-9 to I-84; F-9 to T-83; F-9 to E-82; F-9 to E-81; F-9 to T-80; F-9 to R-79; F-9 to S-78; F-9 to N-77; F-9 to L-76; F-9 to N-75; F-9 to A-74; F-9 to Q-73; F-9 to E-72; F-9 to R-71; F-9 to N-70; F-9 to H-69; F-9 to Q-68; F-9 to W-67; F-9 to G-66; F-9 to G-65; F-9 to K-64; F-9 to R-63; F-9 to L-62; F-9 to Q-61; F-9 to C-60; F-9 to K-59; F-9 to Y-58; F-9 to M-57; F-9 to K-56; F-9 to W-55; F-9 to Y-54; F-9 to E-53; F-9 to P-52; F-9 to Y-51; F-9 to L-50; F-9 to V-49; F-9 to T-48; F-9 to M-47; F-9 to L-46; F-9 to E-45; F-9 to D-44; F-9 to V-43; F-9 to S-42; F-9 to S-41; F-9 to V-40; F-9 to S-39; F-9 to R-38; F-9 to L-37; F-9 to Q-36; F-9 to E-35; F-9 to E-34; F-9 to L-33; F-9 to D-32; F-9 to K-31; F-9 to S-30; F-9 to A-29; F-9 to Y-28; F-9 to A-27; F-9 to T-26; F-9 to A-25; F-9 to E-24; F-9 to G-23; F-9 to A-22; F-9 to D-21; F-9 to P-20; F-9 to E-19; F-9 to A-18; F-9 to D-17; F-9 to S-16; F-9 to L-15; of SEQ ID NO:2. Specifically preferred is the polypeptide fragment comprising amino acid residues F-9 to R-203 of SEQ ID NO:2, as well as polynucleotides encoding this polypeptide. This F-9 to R-203 of SEQ ID NO:2 polypeptide preferably is associated with a S-205 to S-396 of SEQ ID NO:2 polypeptide. Association may be through disulfide, covalent or noncovalent interactions, by linkage via a linker (e.g. serine, glycine, proline linkages), or by an antibody.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:1 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1660 of SEQ ID NO:1, b is an integer of 15 to 1674, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:1, and where the b is greater than or equal to a + 14.

Thus, in one aspect, N-terminal deletion mutants are provided by the present invention. Such mutants include those comprising the amino acid sequence shown in Figure 1 (SEQ ID NO:18) except for a deletion of at least the first 24 N-terminal

amino acid residues (i.e., a deletion of at least Met (1) -- Glu (24)) but not more than the first 115 N-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, first 24 N-terminal amino acid residues (i.e., a deletion of at least Met (1) -- Glu (24)) but not more than the first 103 N-terminal amino acid residues of Figure 1 (SEQ ID NO:18), etc.

In another aspect, C-terminal deletion mutants are provided by the present invention. Such mutants include those comprising the amino acid sequence shown in Figure 1 (SEQ ID NO:18) except for a deletion of at least the last C-terminal amino acid residue (Ser (419)) but not more than the last 220 C-terminal amino acid residues (i.e., a deletion of amino acid residues Val (199) - Ser (419)) of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residue but not more than the last 216 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residue but not more than the last 204 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residues but not more than the last 192 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residues but not more than the last 156 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residues but not more than the last 108 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18). Alternatively, the deletion will include at least the last C-terminal amino acid residues but not more than the last 52 C-terminal amino acid residues of Figure 1 (SEQ ID NO:18).

In yet another aspect, also included by the present invention are deletion mutants having amino acids deleted from both the N-terminal and C-terminal residues. Such mutants include all combinations of the N-terminal deletion mutants and C-terminal deletion mutants described above.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA(s) or the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3 is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and

even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments of 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1025, 1050, 1075, 1100, 1125, 1150, 1175, 1200, 1225, 1250, 1275, 1300, 1325, 1350, 1375, 1400, 1425, 1450, 1475, 1500, 1525, 1550, 1575, 1600, 1625, 1650 or 1674 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the deposited cDNA(s) or as shown in SEQ ID NO:1 or SEQ ID NO:3. By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA(s) or the nucleotide sequence as shown in SEQ ID NOS:1 or 3.

Moreover, representative examples of VEGF-2 polynucleotide fragments include, for example, fragments having a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, or 951 to the end of SEQ ID NO:1 or the cDNA contained in the deposited clone. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity.

Fragments of the full length gene of the present invention may be used as a hybridization probe for a cDNA library to isolate the full length cDNA and to isolate other cDNAs which have a high sequence similarity to the gene or similar biological activity. Probes of this type preferably have at least 30 bases and may contain, for example, 50 or more bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promoter regions, exons, and introns. An example of a screen comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

A VEGF-2 "polynucleotide" also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:1 or for instance, the cDNA clone(s) contained in ATCC Deposit Nos. 97149 or 75698, the complement thereof. "Stringent hybridization conditions" refers to an

overnight incubation at 42° C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

5 Also contemplated are nucleic acid molecules that hybridize to the VEGF-2 polynucleotides at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower
10 stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt
15 concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and
20 commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a
25 complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15
30 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

35 By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the

reference polynucleotide (*e.g.*, the deposited cDNA or the nucleotide sequence as shown in SEQ ID NO:1). Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3N terminal poly(A) tract of the VEGF-2 cDNA shown in SEQ ID NOS:1 or 3), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (*e.g.*, practically any double-stranded cDNA clone).

The present application is directed to nucleic acid molecules at least 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in SEQ ID NOS:1 or 3 or to the nucleic acid sequence of the deposited cDNA(s), irrespective of whether they encode a polypeptide having VEGF-2 activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having VEGF-2 activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having VEGF-2 activity include, *inter alia*, (1) isolating the VEGF-2 gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (*e.g.*, "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the VEGF-2 gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and Northern Blot analysis for detecting VEGF-2 mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence shown in SEQ ID NOS:1 or 3 or to a nucleic acid sequence of the deposited cDNA(s) which do, in fact, encode a polypeptide having VEGF-2 protein activity. By "a polypeptide having VEGF-2 activity" is intended polypeptides exhibiting VEGF-2 activity in a particular biological assay. For example, VEGF-2 protein activity can be measured using, for example, mitogenic assays and endothelial cell migration assays. See, *e.g.*, Olofsson *et al.*, *Proc. Natl. Acad. Sci. USA* 93:2576-2581 (1996) and Joukov *et al.*, *EMBO J.* 5:290-298 (1996).

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence of the deposited cDNA(s) or the nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:3 will encode a polypeptide "having VEGF-2 protein

activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having VEGF-2 protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (*e.g.*, replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

Thus, the present invention is directed to polynucleotides having at least a 70% identity, preferably at least 90% and more preferably at least a 95%, 96%, 97%, or 98% identity to a polynucleotide which encodes the polypeptides of SEQ ID NOS:2 or 4, as well as fragments thereof, which fragments have at least 30 bases and preferably at least 50 bases and to polypeptides encoded by such polynucleotides.

"Identity" *per se* has an art-recognized meaning and can be calculated using published techniques. (See, *e.g.*: (COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, (1988); BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, (1993); COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, (1994); SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, (1987); and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, (1991).) While there exists a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., *SIAM J. Applied Math.* 48:1073 (1988).) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in "Guide to Huge Computers," Martin J. Bishop, ed., Academic Press, San Diego, (1994), and Carillo, H., and Lipton, D., *SIAM J. Applied Math.* 48:1073 (1988). Methods for aligning polynucleotides or polypeptides are codified in computer programs, including the GCG program package (Devereux, J., *et al.*, *Nucleic Acids Research* 12(1):387 (1984)), BLASTP, BLASTN, FASTA (Atschul,

S.F. *et al.*, *J. Molec. Biol.* 215:403 (1990), Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711 (using the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981))). By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the VEGF-2 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence SEQ ID NO:1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag *et al.* (*Comp. App. Biosci.* 6:237-245 (1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent

identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal

positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Table 1 or to the amino acid sequence encoded by deposited DNA clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag *et al.* (*Comp. App. Biosci.* (1990) 6:237-245). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-

terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

VEGF-2 Polypeptides

The present invention further relates to polypeptides which have the deduced amino acid sequence of Figures 1 or 2, or which has the amino acid sequence encoded by the deposited cDNAs, as well as fragments, analogs, and derivatives of such polypeptides.

The terms "fragment," "derivative" and "analog" when referring to the polypeptide of Figures 1 or 2 or that encoded by the deposited cDNA, means a polypeptide which retains the conserved motif of VEGF proteins as shown in Figure 3 and essentially the same biological function or activity.

In the present invention, a "polypeptide fragment" refers to a short amino acid sequence contained in SEQ ID NO:2 or encoded by the cDNA contained in the deposited clone. Protein fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, or 281 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

Preferred polypeptide fragments include the secreted VEGF-2 protein as well as the mature form. Further preferred polypeptide fragments include the secreted VEGF-2 protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted VEGF-2 polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted VEGF-2 protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotide fragments encoding these VEGF-2 polypeptide fragments are also preferred.

Also preferred are VEGF-2 polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:2 falling within conserved domains are specifically contemplated by the present invention. (See Figure 2.) Moreover, polynucleotide fragments encoding these domains are also contemplated.

Other preferred fragments are biologically active VEGF-2 fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the VEGF-2 polypeptide. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

The polypeptides of the present invention may be recombinant polypeptides, natural polypeptides, or synthetic polypeptides, preferably recombinant polypeptides.

It will be recognized in the art that some amino acid sequences of the VEGF-2 polypeptide can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity.

Thus, the invention further includes variations of the VEGF-2 polypeptide which show substantial VEGF-2 polypeptide activity or which include regions of VEGF-2 protein such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent

can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

Thus, the fragments, derivatives, or analogs of the polypeptides of Figures 1 or 2, or that encoded by the deposited cDNAs may be: (I) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code; or (ii) one in which one or more of the amino acid residues includes a substituent group; or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol); or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence; or (v) one in which comprises fewer amino acid residues shown in SEQ ID NOS: 2 or 4, and retains the conserved motif and yet still retains activity characteristics of the VEGF family of polypeptides. Such fragments, derivatives, and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the VEGF-2 protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin. Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845 (1987); Cleland *et al. Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al.*, *Nature* 361:266-268 (1993) describes certain mutations resulting in selective binding of TNF- α to only one of the two known types of TNF receptors. Thus, the VEGF-2 of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

TABLE 1. Conservative Amino Acid Substitutions

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Of course, the number of amino acid substitutions a skilled artisan would make depends on many factors, including those described above. Generally speaking, the number of substitutions for any given VEGF-2 polypeptide will not be more than 50, 40, 30, 25, 20, 15, 10, 5 or 3.

Amino acids in the VEGF-2 protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro*, or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al. Science* 255:306-312 (1992)).

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a

naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

In specific embodiments, the polynucleotides of the invention are less than 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, or 7.5 kb in length. In a further embodiment, polynucleotides of the invention comprise at least 15 contiguous nucleotides of VEGF-2 coding sequence, but do not comprise all or a portion of any VEGF-2 intron. In another embodiment, the nucleic acid comprising VEGF-2 coding sequence does not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the VEGF-2 gene in the genome).

The polypeptides of the present invention include the polypeptides of SEQ ID NOS:2 and 4 (in particular the mature polypeptide) as well as polypeptides which have at least 70% similarity (preferably at least 70% identity) to the polypeptides of SEQ ID NOS:2 and 4, and more preferably at least 90% similarity (more preferably at least 95% identity) to the polypeptides of SEQ ID NOS:2 and 4, and still more preferably at least 95% similarity (still more preferably at least 90% identity) to the polypeptides of SEQ ID NOS:2 and 4 and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide.

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The polypeptides of the present invention include the polypeptide encoded by the deposited cDNA including the leader; the mature polypeptide encoded by the deposited the cDNA minus the leader (i.e., the mature protein); a polypeptide comprising amino acids about - 23 to about 396 in SEQ ID NO:2, a polypeptide comprising amino acids about - 22 to about 396 in SEQ ID NO:2; a polypeptide

comprising amino acids about 1 to about 396 in SEQ ID NO:2; as well as polypeptides which are at least 95% identical, and more preferably at least 96%, 97%, 98% or 99% identical to the polypeptides described above and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

Fusion Proteins

Any VEGF-2 polypeptide can be used to generate fusion proteins. For example, the VEGF-2 polypeptide, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the VEGF-2 polypeptide can be used to indirectly detect the second protein by binding to the VEGF-2. Moreover, because secreted proteins target cellular locations based on trafficking signals, the VEGF-2 polypeptides can be used as a targeting molecule once fused to other proteins.

Examples of domains that can be fused to VEGF-2 polypeptides include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the VEGF-2 polypeptide. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the VEGF-2 polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the VEGF-2 polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the VEGF-2 polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, VEGF-2 polypeptides, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker *et al.*, *Nature* 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis *et al.*, *J. Biochem.* 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D. Bennett *et al.*, *J. Molecular Recognition* 8:52-58 (1995); K. Johanson *et al.*, *J. Biol. Chem.* 270:9459-9471 (1995).)

Moreover, the VEGF-2 polypeptides can be fused to marker sequences, such as a peptide which facilitates purification of VEGF-2. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson *et al.*, *Cell* 37:767 (1984).)

Thus, any of these above fusions can be engineered using the VEGF-2 polynucleotides or the polypeptides.

25 **Biological Activities of VEGF-2**

VEGF-2 polynucleotides and polypeptides can be used in assays to test for one or more biological activities. If VEGF-2 polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that VEGF-2 may be involved in the diseases associated with the biological activity. Therefore, VEGF-2 could be used to treat the associated disease.

30 **Immune Activity**

VEGF-2 polypeptides or polynucleotides may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells,

neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, VEGF-2 polynucleotides or polypeptides can be used as a marker or detector of a particular immune system disease or disorder.

VEGF-2 polynucleotides or polypeptides may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. VEGF-2 polypeptides or polynucleotides could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, VEGF-2 polypeptides or polynucleotides can also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, VEGF-2 polynucleotides or polypeptides could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, VEGF-2 polynucleotides or polypeptides that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting, important in the treatment of heart attacks (infarction), strokes, or scarring.

VEGF-2 polynucleotides or polypeptides may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of VEGF-2 polypeptides or polynucleotides that can inhibit an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by VEGF-2 include, but are not limited to: Addison's Disease, hemolytic anemia,

antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, 5 Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by VEGF-2 10 polypeptides or polynucleotides. Moreover, VEGF-2 can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

VEGF-2 polynucleotides or polypeptides may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs 15 by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of VEGF-2 polypeptides or polynucleotides that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may 20 be an effective therapy in preventing organ rejection or GVHD. Similarly, VEGF-2 polypeptides or polynucleotides may also be used to modulate inflammation. For example, VEGF-2 polypeptides or polynucleotides may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, 25 including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

30

Hyperproliferative Disorders

VEGF-2 polypeptides or polynucleotides can be used to treat or detect hyperproliferative disorders, including neoplasms. VEGF-2 antagonist polypeptides or polynucleotides may inhibit the proliferation of the disorder through direct or 35 indirect interactions. Alternatively, VEGF-2 antagonist polypeptides or

polynucleotides may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Examples of hyperproliferative disorders that can be treated or detected by VEGF-2 antagonist polynucleotides or polypeptides include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by VEGF-2 antagonist polynucleotides or polypeptides. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

Infectious Disease

VEGF-2 polypeptides or polynucleotides can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, VEGF-2 polypeptides or polynucleotides may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by VEGF-2 polynucleotides or polypeptides. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae,

Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. VEGF-2 polypeptides or polynucleotides can be used to treat or detect any of these symptoms or diseases.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by VEGF-2 polynucleotides or polypeptides include, but not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Nocardia), Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (Klebsiella, Salmonella, Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Pasteurellaceae Infections (e.g., Actinobacillus, Haemophilus, Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract

infections, wound infections. VEGF-2 polypeptides or polynucleotides can be used to treat or detect any of these symptoms or diseases.

Moreover, parasitic agents causing disease or symptoms that can be treated or detected by VEGF-2 polynucleotides or polypeptides include, but not limited to, the following families: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas. These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. VEGF-2 polypeptides or polynucleotides can be used to treat or detect any of these symptoms or diseases.

Preferably, treatment using VEGF-2 polypeptides or polynucleotides could either be by administering an effective amount of VEGF-2 polypeptide to the patient, or by removing cells from the patient, supplying the cells with VEGF-2 polynucleotide, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the VEGF-2 polypeptide or polynucleotide can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

VEGF-2 polynucleotides or polypeptides can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, *Science* 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vascular (including vascular endothelium), lymphatic (including lymphatic endothelium), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, VEGF-2 polynucleotides or polypeptides may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. VEGF-2 polynucleotides or polypeptides

of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using VEGF-2 polynucleotides or polypeptides to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the VEGF-2 polynucleotides or polypeptides.

Chemotaxis

VEGF-2 polynucleotides or polypeptides may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

VEGF-2 polynucleotides or polypeptides may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. As a chemotactic molecule, VEGF-2 could also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that VEGF-2 polynucleotides or polypeptides may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, VEGF-2 polynucleotides or polypeptides could be used as an inhibitor of chemotaxis.

Binding Activity

VEGF-2 polypeptides may be used to screen for molecules that bind to VEGF-2 or for molecules to which VEGF-2 binds. The binding of VEGF-2 and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the VEGF-2 or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of VEGF-2, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which VEGF-2 binds (i.e., Flt-4), or at least, a fragment of the receptor capable of being bound by VEGF-2 (e.g., active site). In either case, the molecule can be rationally designed using known techniques. Preferably, the screening for these molecules involves producing appropriate cells which express VEGF-2, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing VEGF-2 (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either VEGF-2 or the molecule.

The assay may simply test binding of a candidate compound to VEGF-2, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to VEGF-2.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing VEGF-2, measuring VEGF-2/molecule activity or binding, and comparing the VEGF-2/molecule activity or binding to a standard.

Preferably, an ELISA assay can measure VEGF-2 level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure VEGF-2 level or activity by either binding, directly or indirectly, to VEGF-2 or by competing with VEGF-2 for a substrate.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the VEGF-2/molecule. Moreover, the assays can discover agents which

may inhibit or enhance the production of VEGF-2 from suitably manipulated cells or tissues.

Therefore, the invention includes a method of identifying compounds which bind to VEGF-2 comprising the steps of: (a) incubating a candidate binding
5 compound with VEGF-2; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with VEGF-2, (b) assaying a biological activity, and (b) determining if a biological activity of VEGF-2 has been altered.

10 Other Activities

VEGF-2 polypeptides or polynucleotides may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

VEGF-2 polypeptides or polynucleotides may also be used to modulate
15 mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, VEGF-2 polypeptides or polynucleotides may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

20 VEGF-2 polypeptides or polynucleotides may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

25 VEGF-2 polypeptides or polynucleotides may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Vectors, Host Cells, and Protein Production

30 The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of VEGF-2 polypeptides or peptides by recombinant techniques.

35 Host cells are genetically engineered (transduced, transformed, or transfected) with the vectors of this invention which may be, for example, a cloning vector or an

expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the VEGF-2 genes of the invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the skilled artisan.

The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide sequence may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, *e.g.*, derivatives of SV40; bacterial plasmids; phage DNA; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other plasmid or vector may be used so long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the *E. coli. lac* or *trp*, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain at least one selectable marker gene to provide a phenotypic trait for selection of transformed host cells. Such markers include dihydrofolate reductase (DHFR) or neomycin resistance for eukaryotic cell culture, and tetracycline or ampicillin resistance for culturing in *E. coli* and other bacteria.

The vector containing the appropriate DNA sequence as herein above described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein. Representative examples of appropriate hosts, include but are not limited to: bacterial cells, such as *E. coli*, *Salmonella typhimurium*, and *Streptomyces*; fungal cells, such

as yeast; insect cells, such as *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, and Bowes melanoma; and plant cells. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

5 More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, 10 for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example - bacterial: pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from 15 Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

20 In addition to the use of expression vectors in the practice of the present invention, the present invention further includes novel expression vectors comprising operator and promoter elements operatively linked to nucleotide sequences encoding a protein of interest. One example of such a vector is pHE4a which is described in detail below.

25 As summarized in Figures 28 and 29, components of the pHE4a vector (SEQ ID NO:16) include: 1) a neomycinphosphotransferase gene as a selection marker, 2) an *E. coli* origin of replication, 3) a T5 phage promoter sequence, 4) two *lac* operator sequences, 5) a Shine-Delgarno sequence, 6) the lactose operon repressor gene (*lacIq*) and 7) a multiple cloning site linker region. The origin of replication (*oriC*) is 30 derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences were made synthetically. Synthetic production of nucleic acid sequences is well known in the art. CLONTECH 95/96 Catalog, pages 215-216, CLONTECH, 1020 East Meadow Circle, Palo Alto, CA 94303. The pHE4a vector was deposited with the ATCC on February 25, 1998, and given accession number 209645.

35 A nucleotide sequence encoding VEGF-2 (SEQ ID NO:1), is operatively linked to the promoter and operator of pHE4a by restricting the vector with NdeI and

either XbaI, BamHI, XhoI, or Asp718, and isolating the larger fragment (the multiple cloning site region is about 310 nucleotides) on a gel. The nucleotide sequence encoding VEGF-2 (SEQ ID NO:1) having the appropriate restriction sites is generated, for example, according to the PCR protocol described in Example 1, using
5 PCR primers having restriction sites for NdeI (as the 5' primer) and either XbaI, BamHI, XhoI, or Asp718 (as the 3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

As noted above, the pHE4a vector contains a *lacIq* gene. *LacIq* is an allele of
10 the *lacI* gene which confers tight regulation of the *lac* operator. Amann, E. *et al.*, *Gene* 69:301-315 (1988); Stark, M., *Gene* 51:255-267 (1987). The *lacIq* gene encodes a repressor protein which binds to *lac* operator sequences and blocks transcription of down-stream (*i.e.*, 3') sequences. However, the *lacIq* gene product dissociates from the *lac* operator in the presence of either lactose or certain lactose
15 analogs, *e.g.*, isopropyl B-D-thiogalactopyranoside (IPTG). VEGF-2 thus is not produced in appreciable quantities in uninduced host cells containing the pHE4a vector. Induction of these host cells by the addition of an agent such as IPTG, however, results in the expression of the VEGF-2 coding sequence.

The promoter/operator sequences of the pHE4a vector (SEQ ID NO:17)
20 comprise a T5 phage promoter and two *lac* operator sequences. One operator is located 5' to the transcriptional start site and the other is located 3' to the same site. These operators, when present in combination with the *lacIq* gene product, confer tight repression of down-stream sequences in the absence of a *lac* operon inducer, *e.g.*, IPTG. Expression of operatively linked sequences located down-stream from
25 the *lac* operators may be induced by the addition of a *lac* operon inducer, such as IPTG. Binding of a *lac* inducer to the *lacIq* proteins results in their release from the *lac* operator sequences and the initiation of transcription of operatively linked sequences. *Lac* operon regulation of gene expression is reviewed in Devlin, T., *TEXTBOOK OF BIOCHEMISTRY WITH CLINICAL CORRELATIONS*, 4th Edition (1997), pages
30 802-807.

The pHE4 series of vectors contain all of the components of the pHE4a vector except for the VEGF-2 coding sequence. Features of the pHE4a vectors include optimized synthetic T5 phage promoter, *lac* operator, and Shine-Delagarno sequences. Further, these sequences are also optimally spaced so that expression of an inserted
35 gene may be tightly regulated and high level of expression occurs upon induction.

Among known bacterial promoters suitable for use in the production of proteins of the present invention include the *E. coli* *lacI* and *lacZ* promoters, the T3 and T7 promoters, the *gpt* promoter, the lambda PR and PL promoters and the *trp* promoter. Suitable eukaryotic promoters include the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous Sarcoma Virus (RSV), and metallothionein promoters, such as the mouse metallothionein-I promoter.

The pHE4a vector also contains a Shine-Delgarno sequence 5' to the AUG initiation codon. Shine-Delgarno sequences are short sequences generally located about 10 nucleotides up-stream (*i.e.*, 5') from the AUG initiation codon. These sequences essentially direct prokaryotic ribosomes to the AUG initiation codon.

Thus, the present invention is also directed to expression vector useful for the production of the proteins of the present invention. This aspect of the invention is exemplified by the pHE4a vector (SEQ ID NO:16).

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include *lacI*, *lacZ*, T3, T7, *gpt*, lambda P_R, P_L and *trp*. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described construct. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, electroporation, transduction, infection, or other methods (Davis, L., *et al.*, *Basic Methods in Molecular Biology* (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with

prokaryotic and eukaryotic hosts are described by Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), the disclosure of which is hereby incorporated by reference.

5 Transcription of a DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin (bp 100 to 270), a cytomegalovirus early
10 promoter enhancer, a polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, *e.g.*, the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a
15 highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), *a-factor*, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader
20 sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a
25 structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E.*
30 *coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication
35 derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for

example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is derepressed by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, well known to those skilled in the art, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, *Cell* 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., VEGF-2 sequence), and/or to include genetic material (e.g., heterologous promoters) that is operably associated with VEGF-2 sequence of the invention, and which activates, alters, and/or amplifies endogenous VEGF-2 polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions and endogenous polynucleotide sequences (e.g. encoding VEGF-2) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature*

342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell; or the host cell
5 can be a prokaryotic cell, such as a bacterial cell. The host strain may be chosen which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide may be controlled. Furthermore, different host
10 cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the protein expressed.

The polypeptides can be recovered and purified from recombinant cell cultures
15 by methods used heretofore, including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. It is preferred to have low concentrations (approximately 0.1-5mM) of calcium ion present during
20 purification (Price *et al.*, *J. Biol. Chem.* 244:917 (1969)). Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides of the present invention may be a naturally purified product, or a
25 product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated with mammalian or other eukaryotic carbohydrates or may be non-
30 glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue.

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, *Proteins: Structures and Molecular Principles*, W.H. Freeman & Co., N.Y., and Hunkapiller, M., *et al.*,
35 1984, *Nature* 310:105-111). For example, a peptide corresponding to a fragment of the VEGF-2 polypeptides of the invention can be synthesized by use of a peptide

synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the VEGF-2 polynucleotide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, γ -methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses VEGF-2 polypeptides which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, *e.g.*, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of VEGF-2 which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U. S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., *Exp. Hematol.* 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (or peptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular

protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

The VEGF-2 polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the VEGF-2 polypeptides of the invention, their preparation, and compositions (preferably, pharmaceutical compositions) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only VEGF-2 polypeptides of the invention (including VEGF-2 fragments, variants, splice variants, and fusion proteins, as described herein). These homomers may contain VEGF-2 polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only VEGF-2 polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing VEGF-2 polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing VEGF-2 polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing VEGF-2 polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the VEGF-2 polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed

when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the VEGF-2 polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in SEQ ID NO:2, or contained in the polypeptide encoded by the deposited clone.) In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a VEGF-2 fusion protein. In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in a VEGF-2-Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another TNF family ligand/receptor member that is capable of forming covalently associated multimers, such as for example, osteoprotegerin (see, e.g., International Publication No. WO 98/49305, the contents of which are herein incorporated by reference in its entirety).

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques

known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Therapeutic Uses

The VEGF-2 polypeptide of the present invention is a potent mitogen for vascular and lymphatic endothelial cells. As shown in Figures 12 and 13, the VEGF-2 polypeptide of SEQ ID NO:2, minus the initial 46 amino acids, is a potent mitogen for vascular endothelial cells and stimulates their growth and proliferation. The results of a Northern blot analysis performed for the VEGF-2 nucleic acid sequence encoding this polypeptide wherein 20 mg of RNA from several human tissues were probed with ³²P-VEGF-2, illustrates that this protein is actively expressed in the heart and lung which is further evidence of mitogenic activity.

Accordingly, VEGF-2, or biologically active portions thereof, may be employed to treat vascular trauma by promoting angiogenesis. For example, to stimulate the growth of transplanted tissue where coronary bypass surgery is performed. VEGF-2, or biologically active portions thereof, may also be employed to promote wound healing, particularly to re-vascularize damaged tissues or stimulate collateral blood flow during ischemia and where new capillary angiogenesis is

desired. VEGF-2, or biologically active portions thereof, may be employed to treat full-thickness wounds such as dermal ulcers, including pressure sores, venous ulcers, and diabetic ulcers. In addition, VEGF-2, or biologically active portions thereof, may be employed to treat full-thickness burns and injuries where a skin graft or flap is used to repair such burns and injuries. VEGF-2, or biologically active portions thereof, may also be employed for use in plastic surgery, for example, for the repair of lacerations, burns, or other trauma. In addition, VEGF-2 can be used to promote healing of wounds and injuries to the eye as well as to treat eye diseases.

Along these same lines, VEGF-2, or biologically active portions thereof, may also be employed to induce the growth of damaged bone, periodontium or ligament tissue. VEGF-2, or biologically active portions thereof, may also be employed for regenerating supporting tissues of the teeth, including cementum and periodontal ligament, that have been damaged by, *e.g.*, periodontal disease or trauma.

Since angiogenesis is important in keeping wounds clean and non-infected, VEGF-2, or biologically active portions thereof, may be employed in association with surgery and following the repair of incisions and cuts. VEGF-2, or biologically active portions thereof, may also be employed for the treatment of abdominal wounds where there is a high risk of infection.

VEGF-2, or biologically active portions thereof, may be employed for the promotion of endothelialization in vascular graft surgery. In the case of vascular grafts using either transplanted or synthetic material, VEGF-2, or biologically active portions thereof, can be applied to the surface of the graft or at the junction to promote the growth of vascular endothelial cells. VEGF-2, or biologically active portions thereof, may also be employed to repair damage of myocardial tissue as a result of myocardial infarction. VEGF-2, or biologically active portions thereof, may also be employed to repair the cardiac vascular system after ischemia. VEGF-2, or biologically active portions thereof, may also be employed to treat damaged vascular tissue as a result of coronary artery disease and peripheral and CNS vascular disease.

VEGF-2, or biologically active portions thereof, may also be employed to coat artificial prostheses or natural organs which are to be transplanted in the body to minimize rejection of the transplanted material and to stimulate vascularization of the transplanted materials.

VEGF-2, or biologically active portions thereof, may also be employed for vascular tissue repair of injuries resulting from trauma, for example, that occurring during arteriosclerosis and required following balloon angioplasty where vascular tissues are damaged.

VEGF-2, or biologically active portions thereof, may also be used to treat peripheral arterial disease. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat peripheral arterial disease. Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of
5 alleviating or treating peripheral arterial disease. Suitable doses, formulations, and administration routes are described below.

VEGF-2, or biologically active portions thereof, may also to promote the endothelial function of lymphatic tissues and vessels, such as to treat the loss of lymphatic vessels, occlusions of lymphatic vessels, and lymphangiomas. VEGF-2
10 may also be used to stimulate lymphocyte production.

VEGF-2, or biologically active portions thereof, may also be used to treat hemangioma in newborns. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat hemangioma in newborns. Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of
15 alleviating or treating hemangioma in newborns. Suitable doses, formulations, and administration routes are described below.

VEGF-2, or biologically active portions thereof, may also be used to prevent or treat abnormal retinal development in premature newborns. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat
20 abnormal retinal development in premature newborns. Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of alleviating or treating abnormal retinal development in premature newborns. Suitable doses, formulations, and administration routes are described below.

VEGF-2, or biologically active portions thereof, may be used to treat primary
25 (idiopathic) lymphademas, including Milroy's disease and Lymphedema praecox. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat primary (idiopathic) lymphademas, including Milroy's disease and Lymphedema praecox. Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of alleviating or treating primary (idiopathic) lymphademas,
30 including Milroy's disease and Lymphedema praecox. VEGF-2 or biologically active portions thereof, may also be used to treat edema as well as to effect blood pressure in an animal. Suitable doses, formulations, and administration routes are described below.

VEGF-2, or biologically active portions thereof, may also be used to treat
35 secondary (obstructive) lifetimes including those that result from (i) the removal of lymph nodes and vessels, (ii) radiotherapy and surgery in the treatment of cancer, and

(iii) trauma and infection. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat secondary (obstructive) lifetimes including those that result from (I) the removal of lymph nodes and vessels, (ii) radiotherapy and surgery in the treatment of cancer, and (iii) trauma and infection.

5 Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of secondary (obstructive) lifetimes including those that result from (I) the removal of lymph nodes and vessels, (ii) radiotherapy and surgery in the treatment of cancer, and (iii) trauma and infection. Suitable doses, formulations, and administration routes are described below.

10 VEGF-2, or biologically active portions thereof, may also be used to treat Kaposi's Sarcoma. Accordingly, in a further aspect, there is provided a process for utilizing VEGF-2 polypeptides to treat Kaposi's Sarcoma. Preferably, a VEGF-2 polypeptide is administered to an individual for the purpose of alleviating or treating Kaposi's Sarcoma. Suitable doses, formulations, and administration routes are
15 described below.

VEGF-2 antagonists can be used to treat cancer by inhibiting the angiogenesis necessary to support cancer and tumor growth.

Cardiovascular Disorders

20 The present inventors have shown that VEGF-2 stimulates the growth of vascular endothelial cells, stimulates endothelial cell migration, stimulates angiogenesis in the CAM assay, decreases blood pressure in spontaneously hypertensive rats, and increases blood flow to ischemic limbs in rabbits. Accordingly, VEGF-2 polypeptides or polynucleotides encoding VEGF-2 may be
25 used to treat cardiovascular disorders, including peripheral artery disease, such as limb ischemia.

Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects
30 include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion
35 defects, Lutembacher's Syndrome, trilogly of Fallot, ventricular heart septal defects.

Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, angiodyspiasia, angiomatosis, bacillary angiomatosis, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic

edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, atacia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subaraxhnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

VEGF-2 polypeptides or polynucleotides are especially effective for the treatment of critical-limb ischemia and coronary disease. As shown in Example 18, administration of VEGF-2 polynucleotides and polypeptides to an experimentally induced ischemia rabbit hindlimb restored blood pressure ratio, blood flow, angiographic score, and capillary density.

VEGF-2 polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. VEGF-2 polypeptides may be administered as part of a pharmaceutical composition, described in more detail below. Methods of delivering VEGF-2 polynucleotides are described in more detail below.

15.

Gene Therapy Methods

Another aspect of the present invention is to gene therapy methods for treating disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of the VEGF-2 polypeptide of the present invention. This method requires a polynucleotide which codes for a VEGF-2 polypeptide operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO 90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a VEGF-2 polynucleotide *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Beldegrun, A., *et al.*, *J. Natl. Cancer Inst.* 85: 207-216 (1993); Ferrantini, M. *et al.*, *Cancer Research* 53: 1107-1112 (1993); Ferrantini, M. *et al.*, *J. Immunology* 153: 4604-4615 (1994); Kaido, T., *et al.*, *Int. J. Cancer* 60: 221-229 (1995); Ogura, H., *et al.*, *Cancer Research* 50: 5102-5106 (1990); Santodonato, L., *et al.*, *Human Gene Therapy* 7:1-10 (1996); Santodonato, L., *et al.*, *Gene Therapy* 4:1246-1255 (1997); and Zhang, J.-F. *et al.*, *Cancer Gene Therapy* 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered

are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the VEGF-2 polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The VEGF-2 polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the VEGF-2 polynucleotide is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the VEGF-2 polynucleotides can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

The VEGF-2 polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of VEGF-2 DNA. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for VEGF-2.

Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA

sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The VEGF-2 polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked VEGF-2 DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

As is evidenced by Example 18, naked VEGF-2 nucleic acid sequences can be administered *in vivo* results in the successful expression of VEGF-2 polypeptide in the femoral arteries of rabbits.

The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

In certain embodiments, the VEGF-2 polynucleotide constructs are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* (1987) 84:7413-7416, which is herein incorporated by reference); mRNA (Malone *et al.*, *Proc. Natl. Acad. Sci. USA* (1989) 86:6077-6081, which is herein incorporated by reference); and purified transcription factors (Debs *et al.*, *J. Biol. Chem.* (1990) 265:10189-10192, which is herein incorporated by reference), in functional form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* (1987) 84:7413-7416, which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication No. WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., P. Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE),

among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC),
5 dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is
10 hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to
15 produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known
20 in the art. See, e.g., Straubinger *et al.*, *Methods of Immunology* (1983), 101:512-527, which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a
25 homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The
30 liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos *et al.*, *Biochim. Biophys. Acta* (1975) 394:483; Wilson *et al.*, *Cell* (1979) 17:77); ether
35 injection (Deamer, D. and Bangham, A., *Biochim. Biophys. Acta* (1976) 443:629, Ostro *et al.*, *Biochem. Biophys. Res. Commun.* (1977) 76:836; Fraley *et al.*, *Proc.*

Natl. Acad. Sci. USA (1979) 76:3348); detergent dialysis (Enoch, H. and Strittmatter, P., *Proc. Natl. Acad. Sci. USA* (1979) 76:145); and reverse-phase evaporation (REV) (Fraley *et al.*, *J. Biol. Chem.* (1980) 255:10431; Szoka, F. and Papahadjopoulos, D., *Proc. Natl. Acad. Sci. USA* (1978) 75:145; Schaefer-Ridder *et al.*, *Science* (1982) 215:166), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent No. 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are be engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding VEGF-2. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, *Human Gene Therapy* 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding VEGF-2. Such retroviral vector particles then may

be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express VEGF-2.

In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with VEGF-2 polynucleotide contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses VEGF-2, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz, A. R. *et al.* (1974) *Am. Rev. Respir. Dis.* 109:233-238). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld, M. A. *et al.* (1991) *Science* 252:431-434; Rosenfeld *et al.*, (1992) *Cell* 68:143-155). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green, M. *et al.* (1979) *Proc. Natl. Acad. Sci. USA* 76:6606).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, *Curr. Opin. Genet. Devel.* 3:499-503 (1993); Rosenfeld *et al.*, *Cell* 68:143-155 (1992); Engelhardt *et al.*, *Human Genet. Ther.* 4:759-769 (1993); Yang *et al.*, *Nature Genet.* 7:362-369 (1994); Wilson *et al.*, *Nature* 365:691-692 (1993); and U.S. Patent No. 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express E1a and E1b, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, for example, the HARP promoter of the present invention, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, N., *Curr. Topics in Microbiol. Immunol.* 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The VEGF-2 polynucleotide construct is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the VEGF-2 polynucleotide construct. These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the VEGF-2 polynucleotide construct integrated into its genome, and will express VEGF-2.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding VEGF-2) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature* 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of

the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the VEGF-2 desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

5 The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the
10 amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating
15 agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous
20 recombination between the construct and the endogenous sequence takes place, such that an endogenous VEGF-2 sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous VEGF-2 sequence.

The polynucleotides encoding VEGF-2 may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include,
25 but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

30 Preferably, the polynucleotide encoding VEGF-2 contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells
35 to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers (Kaneda *et al.*, *Science* 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling *et al.*, *Proc. Natl. Acad. Sci. USA* 189:11277-11281, 1992, which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a

polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly preferred.

Nucleic Acid Utilities

VEGF-2 nucleic acid sequences and VEGF-2 polypeptides may also be employed for *in vitro* purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors, and for the production of diagnostics and therapeutics to treat human disease. For example, VEGF-2 may be employed for *in vitro* culturing of vascular endothelial cells, where it is added to the conditional medium in a concentration from 10 pg/ml to 10 ng/ml.

Fragments of the full length VEGF-2 gene may be used as a hybridization probe for a cDNA library to isolate other genes which have a high sequence similarity to the gene or similar biological activity. Probes of this type generally have at least 50 base pairs, although they may have a greater number of bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete VEGF-2 gene including regulatory and promoter regions, exons, and introns. An example of a screen comprises isolating the coding region of the VEGF-2 gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

This invention provides methods for identification of VEGF-2 receptors. The gene encoding the receptor can be identified by numerous methods known to those of

skill in the art, for example, ligand panning and FACS sorting (Coligan *et al.*, *Current Protocols in Immun.*, 1(2), Chapter 5, (1991)). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to VEGF-2, and a cDNA library created from this RNA is divided into pools and used to
5 transfect COS cells or other cells that are not responsive to VEGF-2. Transfected cells which are grown on glass slides are exposed to labeled VEGF-2. VEGF-2 can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are
10 prepared and retransfected using an iterative sub-pooling and rescreening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, labeled VEGF-2 can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray
15 film. The labeled complex containing VEGF-2 is then excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

20 *VEGF-2 Agonist and Antagonists*

This invention is also related to a method of screening compounds to identify those which are VEGF-2 agonists or antagonists. An example of such a method takes advantage of the ability of VEGF-2 to significantly stimulate the proliferation of
25 human endothelial cells in the presence of the mitogen Con A. Endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) in a reaction mixture supplemented with Con-A (Calbiochem, La Jolla, CA). Con-A, polypeptides of the present invention and the compound to be screened are added. After incubation at 37°C, cultures are pulsed with 1 FCi of ^3H -thymidine (5
30 Ci/mmol; 1 Ci = 37 BGq; NEN) for a sufficient time to incorporate the ^3H and harvested onto glass fiber filters (Cambridge Technology, Watertown, MA). Mean ^3H -thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant ^3H -thymidine
35 incorporation, as compared to a control assay where the compound is excluded, indicates stimulation of endothelial cell proliferation.

To assay for antagonists, the assay described above is performed and the ability of the compound to inhibit ³[H]thymidine incorporation in the presence of VEGF-2 indicates that the compound is an antagonist to VEGF-2. Alternatively, VEGF-2 antagonists may be detected by combining VEGF-2 and a potential antagonist with membrane-bound VEGF-2 receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. VEGF-2 can be labeled, such as by radioactivity, such that the number of VEGF-2 molecules bound to the receptor can determine the effectiveness of the potential antagonist.

Alternatively, the response of a known second messenger system following interaction of VEGF-2 and receptor would be measured and compared in the presence or absence of the compound. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis. In another method, a mammalian cell or membrane preparation expressing the VEGF-2 receptor is incubated with labeled VEGF-2 in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured.

Potential VEGF-2 antagonists include an antibody, or in some cases, an oligonucleotide, which bind to the polypeptide and effectively eliminate VEGF-2 function. Alternatively, a potential antagonist may be a closely related protein which binds to VEGF-2 receptors, however, they are inactive forms of the polypeptide and thereby prevent the action of VEGF-2. Examples of these antagonists include a negative dominant mutant of the VEGF-2 polypeptide, for example, one chain of the hetero-dimeric form of VEGF-2 may be dominant and may be mutated such that biological activity is not retained. An example of a negative dominant mutant includes truncated versions of a dimeric VEGF-2 which is capable of interacting with another dimer to form wild type VEGF-2, however, the resulting homo-dimer is inactive and fails to exhibit characteristic VEGF activity.

Another potential VEGF-2 antagonist is an antisense construct prepared using antisense technology. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes for the mature polypeptides of the present invention, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix -see Lee *et al.*, *Nucl. Acids Res.* 6:3073 (1979); Cooney *et al.*, *Science* 241:456 (1988); and Dervan *et al.*, *Science* 251:1360 (1991)), thereby preventing transcription and the

production of VEGF-2. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the VEGF-2 polypeptide (Antisense - Okano, *J. Neurochem.* 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). The
5 oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of VEGF-2.

Potential VEGF-2 antagonists also include small molecules which bind to and occupy the active site of the polypeptide thereby making the catalytic site inaccessible to substrate such that normal biological activity is prevented. Examples of small
10 molecules include but are not limited to small peptides or peptide-like molecules.

The antagonists may be employed to limit angiogenesis necessary for solid tumor metastasis. The identification of VEGF-2 can be used for the generation of certain inhibitors of vascular endothelial growth factor. Since angiogenesis and neovascularization are essential steps in solid tumor growth, inhibition of angiogenic
15 activity of the vascular endothelial growth factor is very useful to prevent the further growth, retard, or even regress solid tumors. Although the level of expression of VEGF-2 is extremely low in normal tissues including breast, it can be found expressed at moderate levels in at least two breast tumor cell lines that are derived from malignant tumors. It is, therefore, possible that VEGF-2 is involved in tumor
20 angiogenesis and growth.

Gliomas are also a type of neoplasia which may be treated with the antagonists of the present invention.

The antagonists may also be used to treat chronic inflammation caused by increased vascular permeability. In addition to these disorders, the antagonists may
25 also be employed to treat retinopathy associated with diabetes, rheumatoid arthritis and psoriasis.

The antagonists may be employed in a composition with a pharmaceutically acceptable carrier, *e.g.*, as hereinafter described.

30 ***Pharmaceutical compositions***

The VEGF-2 polypeptides, polynucleotides and agonists and antagonists may be employed in combination with a suitable pharmaceutical carrier. Such compositions comprise a therapeutically effective amount of the polypeptide or agonist or antagonist, and a pharmaceutically acceptable carrier or excipient. Such a carrier
35 includes but is not limited to saline, buffered saline, dextrose, water, glycerol,

ethanol, and combinations thereof. The formulation should suit the mode of administration.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the pharmaceutical compositions may be employed in conjunction with other therapeutic compounds.

The pharmaceutical compositions may be administered in a convenient manner such as by the topical, intravenous, intraperitoneal, intramuscular, intratumor, subcutaneous, intranasal or intradermal routes. The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In general, the pharmaceutical compositions are administered in an amount of at least about 10 mg/kg body weight and in most cases they will be administered in an amount not in excess of about 8 mg/Kg body weight per day. In most cases, the dosage is from about 10 mg/kg to about 1 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc.

The VEGF-2 polypeptides, and agonists or antagonists which are polypeptides may also be employed in accordance with the present invention by expression of such polypeptide *in vivo*, which is often referred to as "gene therapy," described above.

Thus, for example, cells such as bone marrow cells may be engineered with a polynucleotide (DNA or RNA) encoding for the polypeptide *ex vivo*, the engineered cells are then provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding the polypeptide of the present invention.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo*, for example, by procedures known in the art. As known in the art, a producer cell for producing a retroviral particle containing RNA encoding a polypeptide of the present invention may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such methods should be apparent to those skilled in the art from the teachings of the present invention. For example, the expression vehicle for engineering cells may be other than a retroviral particle, for

example, an adenovirus, which may be used to engineer cells *in vivo* after combination with a suitable delivery vehicle.

Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

The vector includes one or more promoters. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller *et al.*, *Biotechniques* 7:980-990 (1989), or any other promoter (*e.g.*, cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and b-actin promoters). Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

The nucleic acid sequence encoding the polypeptide of the present invention is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs hereinabove described); the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter which controls the gene encoding the polypeptide.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, y-2, y-AM, PA12, T19-14X, VT-19-17-H2, yCRE, yCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, *Human Gene Therapy* 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to,

electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

Diagnostic assays

This invention is also related to the use of the VEGF-2 gene as part of a diagnostic assay for detecting diseases or susceptibility to diseases related to the presence of mutations in VEGF-2 nucleic acid sequences.

Individuals carrying mutations in the VEGF-2 gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki *et al.*, *Nature* 324:163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding VEGF-2 can be used to identify and analyze VEGF-2 mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled VEGF-2 RNA or alternatively, radiolabeled VEGF-2 antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their

specific melting or partial melting temperatures (see, *e.g.*, Myers *et al.*, *Science* 230:1242 (1985)).

Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (*e.g.*, Cotton *et al.*, *PNAS, USA* 85:4397-4401 (1985)).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (*e.g.*, Restriction Fragment Length Polymorphisms (RFLP)) and Southern blotting of genomic DNA.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

The present invention also relates to a diagnostic assay for detecting altered levels of VEGF-2 protein in various tissues since an over-expression of the proteins compared to normal control tissue samples may detect the presence of a disease or susceptibility to a disease, for example, abnormal cellular differentiation. Assays used to detect levels of VEGF-2 protein in a sample derived from a host are well-known to those of skill in the art and include radioimmunoassays, competitive-binding assays, Western Blot analysis, ELISA assays and "sandwich" assay. An ELISA assay (Coligan *et al.*, *Current Protocols in Immunology* 1(2), Chapter 6, (1991)) initially comprises preparing an antibody specific to the VEGF-2 antigen, preferably a monoclonal antibody. In addition a reporter antibody is prepared against the monoclonal antibody. To the reporter antibody is attached a detectable reagent such as radioactivity, fluorescence or, in this example, a horseradish peroxidase enzyme. A sample is removed from a host and incubated on a solid support, *e.g.* a polystyrene dish, that binds the proteins in the sample. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein, such as, bovine serum albumen. Next, the monoclonal antibody is incubated in the dish during which time the monoclonal antibodies attach to any VEGF-2 proteins attached to the polystyrene dish. All unbound monoclonal antibody is washed out with buffer. The reporter antibody linked to horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to VEGF-2. Unattached reporter antibody is then washed out. Peroxidase substrates are then added to the dish and the amount of color developed in a given time period is a measurement of the amount of VEGF-2 protein present in a given volume of patient sample when compared against a standard curve.

A competition assay may be employed wherein antibodies specific to VEGF-2 are attached to a solid support. Polypeptides of the present invention are then labeled, for example, by radioactivity, and a sample derived from the host are passed over the solid support and the amount of label detected, for example by liquid scintillation chromatography, can be correlated to a quantity of VEGF-2 in the sample.

A "sandwich" assay is similar to an ELISA assay. In a "sandwich" assay VEGF-2 is passed over a solid support and binds to antibody attached to a solid support. A second antibody is then bound to the VEGF-2. A third antibody which is labeled and specific to the second antibody is then passed over the solid support and binds to the second antibody and an amount can then be quantified.

Chromosome identification

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphism's) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the cDNA is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one

step. This technique can be used with probes from the cDNA as short as 50 or 60 base pairs. For a review of this technique, see Verma *et al.*, *Human Chromosomes: a Manual of Basic Techniques*, Pergamon Press, New York (1988).

5 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of
10 physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

15 With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

Comparison of affected and unaffected individuals generally involves first
20 looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that cDNA sequence. Ultimately, complete sequencing of genes from several individuals is required to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

25

Antisense

The present invention is further directed to inhibiting VEGF-2 *in vivo* by the use of antisense technology. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which
30 methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the mature polynucleotide sequence, which encodes for the polypeptide of the present invention, is used to design an antisense RNA oligonucleotide of from 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple
35 helix -- see Lee *et al.*, *Nucl. Acids Res.* 6:3073 (1979); Cooney *et al.*, *Science* 241:456 (1988); and Dervan *et al.* *Science*, 251:1360 (1991), thereby preventing

transcription and the production of VEGF-2. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of an mRNA molecule into the VEGF-2 (antisense -- Okano, *J. Neurochem.* 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)).

- 5 Alternatively, the oligonucleotides described above can be delivered to cells by procedures in the art such that the anti-sense RNA or DNA may be expressed *in vivo* to inhibit production of VEGF-2 in the manner described above.

Antisense constructs to VEGF-2, therefore, may inhibit the angiogenic activity of the VEGF-2 and prevent the further growth or even regress solid tumors, since
10 angiogenesis and neovascularization are essential steps in solid tumor growth. These antisense constructs may also be used to treat rheumatoid arthritis, psoriasis, diabetic retinopathy and Kaposi's sarcoma which are all characterized by abnormal angiogenesis.

15 ***Epitope-Bearing Portions***

In another aspect, the invention provides peptides and polypeptides comprising epitope-bearing portions of the polypeptides of the present invention. These epitopes are immunogenic or antigenic epitopes of the polypeptides of the present invention. An "immunogenic epitope" is defined as a part of a protein that
20 elicits an antibody response *in vivo* when the whole polypeptide of the present invention, or fragment thereof, is the immunogen. On the other hand, a region of a polypeptide to which an antibody can bind is defined as an "antigenic determinant" or "antigenic epitope." The number of *in vivo* immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, e.g., Geysen, et al.
25 (1983) Proc. Natl. Acad. Sci. USA 81:3998-4002. However, antibodies can be made to any antigenic epitope, regardless of whether it is an immunogenic epitope, by using methods such as phage display. See e.g., Petersen G. et al. (1995) Mol. Gen. Genet. 249:425-431. Therefore, included in the present invention are both immunogenic epitopes and antigenic epitopes.

30 It is particularly pointed out that the immunogenic epitopes comprises predicted critical amino acid residues determined by the Jameson-Wolf analysis. Thus, additional flanking residues on either the N-terminal, C-terminal, or both N- and C-terminal ends may be added to these sequences to generate an epitope-bearing polypeptide of the present invention. Therefore, the immunogenic epitopes may
35 include additional N-terminal or C-terminal amino acid residues. The additional flanking amino acid residues may be contiguous flanking N-terminal and/or C-

terminal sequences from the polypeptides of the present invention, heterologous polypeptide sequences, or may include both contiguous flanking sequences from the polypeptides of the present invention and heterologous polypeptide sequences.

Polypeptides of the present invention comprising immunogenic or antigenic epitopes are at least 7 amino acids residues in length. "At least" means that a polypeptide of the present invention comprising an immunogenic or antigenic epitope may be 7 amino acid residues in length or any integer between 7 amino acids and the number of amino acid residues of the full length polypeptides of the invention. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. However, it is pointed out that each and every integer between 7 and the number of amino acid residues of the full length polypeptide are included in the present invention.

The immuno and antigenic epitope-bearing fragments may be specified by either the number of contiguous amino acid residues, as described above, or further specified by N-terminal and C-terminal positions of these fragments on the amino acid sequence of SEQ ID NO:2. Every combination of a N-terminal and C-terminal position that a fragment of, for example, at least 7 or at least 15 contiguous amino acid residues in length could occupy on the amino acid sequence of SEQ ID NO:2 is included in the invention. Again, "at least 7 contiguous amino acid residues in length" means 7 amino acid residues in length or any integer between 7 amino acids and the number of amino acid residues of the full length polypeptide of the present invention. Specifically, each and every integer between 7 and the number of amino acid residues of the full length polypeptide are included in the present invention.

Immunogenic and antigenic epitope-bearing polypeptides of the invention are useful, for example, to make antibodies which specifically bind the polypeptides of the invention, and in immunoassays to detect the polypeptides of the present invention. The antibodies are useful, for example, in affinity purification of the polypeptides of the present invention. The antibodies may also routinely be used in a variety of qualitative or quantitative immunoassays, specifically for the polypeptides of the present invention using methods known in the art. *See, e.g.,* Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press; 2nd Ed. 1988).

The epitope-bearing polypeptides of the present invention may be produced by any conventional means for making polypeptides including synthetic and recombinant methods known in the art. For instance, epitope-bearing peptides may be synthesized

using known methods of chemical synthesis. For instance, Houghten has described a simple method for the synthesis of large numbers of peptides, such as 10-20-mgs of 248 individual and distinct 13 residue peptides representing single amino acid variants of a segment of the HA1 polypeptide, all of which were prepared and characterized (by ELISA-type binding studies) in less than four weeks (Houghten, R. A. Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985)). This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten and coworkers (1986). In this procedure the individual resins for the solid-phase synthesis of various peptides are contained in separate solvent-permeable packets, enabling the optimal use of the many identical repetitive steps involved in solid-phase methods. A completely manual procedure allows 500-1000 or more syntheses to be conducted simultaneously (Houghten et al. (1985) Proc. Natl. Acad. Sci. 82:5131-5135 at 5134).

Epitope-bearing polypeptides of the present invention are used to induce antibodies according to methods well known in the art including, but not limited to, *in vivo* immunization, *in vitro* immunization, and phage display methods. See, e.g., Sutcliffe, et al., *supra*; Wilson, et al., *supra*, and Bittle, et al. (1985) J. Gen. Virol. 66:2347-2354. If *in vivo* immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling of the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as -maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier-coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µgs of peptide or carrier protein and Freund's adjuvant. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to heterologous polypeptide sequences. For example, the polypeptides of the present

invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, any combination thereof including both entire domains and portions thereof) resulting in chimeric polypeptides. These fusion proteins facilitate purification, and show an increased half-life *in vivo*. This has been shown, *e.g.*, for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. *See, e.g.*, EPA 0,394,827; Traunecker et al. (1988) Nature 331:84-86. Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion can also be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. *See, e.g.*, Fountoulakis et al. (1995) J. Biochem. 270:3958-3964. Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag to aid in detection and purification of the expressed polypeptide.

15 **Antibodies**

The present invention further relates to antibodies and T-cell antigen receptors (TCR) which specifically bind the polypeptides of the present invention. The antibodies of the present invention include IgG (including IgG1, IgG2, IgG3, and IgG4), IgA (including IgA1 and IgA2), IgD, IgE, or IgM, and IgY. As used herein, the term "antibody" (Ab) is meant to include whole antibodies, including single-chain whole antibodies, and antigen-binding fragments thereof. Most preferably the antibodies are human antigen binding antibody fragments of the present invention include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a V_L or V_H domain. The antibodies may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, rabbit, goat, guinea pig, camel, horse, or chicken.

Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entire or partial of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are any combinations of variable region(s) and hinge region, CH1, CH2, and CH3 domains. The present invention further includes chimeric, humanized, and human monoclonal and polyclonal antibodies which specifically bind the polypeptides of the present invention. The present invention further includes antibodies which are anti-idiotypic to the antibodies of the present invention.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for heterologous compositions, such as a heterologous polypeptide or solid support material. *See, e.g.*, WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, A. et al. (1991) J. Immunol. 147:60-69; US Patents 5,573,920, 4,474,893, 5,601,819, 4,714,681, 4,925,648; Kostelny, S.A. et al. (1992) J. Immunol. 148:1547-1553.

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which are recognized or specifically bound by the antibody. The epitope(s) or polypeptide portion(s) may be specified as described herein, *e.g.*, by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of the polypeptides of the present invention are included. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. Further included in the present invention are antibodies which only bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity. Preferred binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$, $10^{-8}M$, $5 \times 10^{-9}M$, $10^{-9}M$, $5 \times 10^{-10}M$, $10^{-10}M$, $5 \times 10^{-11}M$, $10^{-11}M$, $5 \times 10^{-12}M$, $10^{-12}M$, $5 \times 10^{-13}M$, $10^{-13}M$, $5 \times 10^{-14}M$, $10^{-14}M$, $5 \times 10^{-15}M$, and $10^{-15}M$.

Antibodies of the present invention have uses that include, but are not limited to, methods known in the art to purify, detect, and target the polypeptides of the present invention including both *in vitro* and *in vivo* diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and

quantitatively measuring levels of the polypeptides of the present invention in biological samples. *See, e.g.*, Harlow et al., *ANTIBODIES: A LABORATORY MANUAL*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference in the entirety).

5 The antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly
10 fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, or toxins. *See, e.g.*, WO 92/08495; WO 91/14438; WO 89/12624; US Patent 5,314,995; and EP 0 396 387. The antibodies of the present invention may be prepared by any suitable method known in the art. For example, a polypeptide of the present invention or an antigenic
15 fragment thereof can be administered to an animal in order to induce the production of sera containing polyclonal antibodies. Monoclonal antibodies can be prepared using a wide of techniques known in the art including the use of hybridoma and recombinant technology. *See, e.g.*, Harlow et al., *ANTIBODIES: A LABORATORY MANUAL*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in:
20 *MONOCLONAL ANTIBODIES AND T-CELL HYBRIDOMAS* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). Fab and F(ab')₂ fragments may be produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments).

Alternatively, antibodies of the present invention can be produced through the
25 application of recombinant DNA technology or through synthetic chemistry using methods known in the art. For example, the antibodies of the present invention can be prepared using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of a phage particle which carries polynucleotide sequences encoding them. Phage with a desired binding
30 property are selected from a repertoire or combinatorial antibody library (e.g. human or murine) by selecting directly with antigen, typically antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of
35 phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman U. et al. (1995) *J. Immunol. Methods* 182:41-

50; Ames, R.S. et al. (1995) J. Immunol. Methods 184:177-186; Kettleborough, C.A. et al. (1994) Eur. J. Immunol. 24:952-958; Persic, L. et al. (1997) Gene 187 9-18; Burton, D.R. et al. (1994) Advances in Immunology 57:191-280; PCT/GB91/01134; WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and US Patents 5,698,426, 5,223,409, 5,403,484, 5,580,717, 5,427,908, 5,750,753, 5,821,047, 5,571,698, 5,427,908, 5,516,637, 5,780,225, 5,658,727 and 5,733,743 (said references incorporated by reference in their entirety).

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host including mammalian cells, insect cells, plant cells, yeast, and bacteria. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in WO 92/22324; Mullinax, R.L. et al. (1992) BioTechniques 12(6):864-869; and Sawai, H. et al. (1995) AJRI 34:26-34; and Better, M. et al. (1988) Science 240:1041-1043 (said references incorporated by reference in their entirety).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al. (1991) Methods in Enzymology 203:46-88; Shu, L. et al. (1993) PNAS 90:7995-7999; and Skerra, A. et al. (1988) Science 240:1038-1040. For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies, S.D. et al. (1989) J. Immunol. Methods 125:191-202; and US Patent 5,807,715. Antibodies can be humanized using a variety of techniques including CDR-grafting (EP 0 239 400; WO 91/09967; US Patent 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan E.A., (1991) Molecular Immunology 28(4/5):489-498; Studnicka G.M. et al. (1994) Protein Engineering 7(6):805-814; Roguska M.A. et al. (1994) PNAS 91:969-973), and chain shuffling (US Patent 5,565,332). Human antibodies can be made by a variety of methods known in the art including phage display methods described above. See also, US Patents 4,444,887, 4,716,111, 5,545,806, and 5,814,318; and WO 98/46645 (said references incorporated by reference in their entirety).

Further included in the present invention are antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide of the present invention. The antibodies may be specific for antigens other than polypeptides of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either *in vitro* or *in vivo*, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in *in vitro* immunoassays and purification methods using methods known in the art. *See e.g.*, Harbor et al. *supra* and WO 93/21232; EP 0 439 095; Naramura, M. et al. (1994) Immunol. Lett. 39:91-99; US Patent 5,474,981; Gillies, S.O. et al. (1992) PNAS 89:1428-1432; Fell, H.P. et al. (1991) J. Immunol. 146:2446-2452 (said references incorporated by reference in their entireties).

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides of the present invention may be fused or conjugated to the above antibody portions to increase the *in vivo* half life of the polypeptides or for use in immunoassays using methods known in the art. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. *See e.g.*, US Patents 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, 5,112,946; EP 0 307 434, EP 0 367 166; WO 96/04388, WO 91/06570; Ashkenazi, A. et al. (1991) PNAS 88:10535-10539; Zheng, X.X. et al. (1995) J. Immunol. 154:5590-5600; and Vil. H. et al. (1992) PNAS 89:11337-11341 (said references incorporated by reference in their entireties).

The invention further relates to antibodies which act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the

polypeptides of the invention either partially or fully. Included are both receptor-specific antibodies and ligand-specific antibodies. Included are receptor-specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. Also include are receptor-specific antibodies which both prevent ligand binding and receptor activation. Likewise, included are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included are antibodies which activate the receptor. These antibodies may act as agonists for either all or less than all of the biological activities affected by ligand-mediated receptor activation. The antibodies may be specified as agonists or antagonists for biological activities comprising specific activities disclosed herein. The above antibody agonists can be made using methods known in the art. See e.g., WO 96/40281; US Patent 5,811,097; Deng, B. et al. (1998) *Blood* 92(6):1981-1988; Chen, Z. et al. (1998) *Cancer Res.* 58(16):3668-3678; Harrop, J.A. et al. (1998) *J. Immunol.* 161(4):1786-1794; Zhu, Z. et al. (1998) *Cancer Res.* 58(15):3209-3214; Yoon, D.Y. et al. (1998) *J. Immunol.* 160(7):3170-3179; Prat, M. et al. (1998) *J. Cell. Sci.* 111(Pt2):237-247; Pitard, V. et al. (1997) *J. Immunol. Methods* 205(2):177-190; Liautard, J. et al. (1997) *Cytokine* 9(4):233-241; Carlson, N.G. et al. (1997) *J. Biol. Chem.* 272(17):11295-11301; Taryman, R.E. et al. (1995) *Neuron* 14(4):755-762; Muller, Y.A. et al. (1998) *Structure* 6(9):1153-1167; Bartunek, P. et al. (1996) *Cytokine* 8(1):14-20 (said references incorporated by reference in their entireties).

Antibodies may further be used in an immunoassay to detect the presence of tumors in certain individuals. Enzyme immunoassay can be performed from the blood sample of an individual. Elevated levels of VEGF-2 can be considered diagnostic of cancer.

Truncated versions of VEGF-2 can also be produced that are capable of interacting with wild type VEGF-2 to form dimers that fail to activate endothelial cell growth, therefore inactivating the endogenous VEGF-2. Or, mutant forms of VEGF-2 form dimers themselves and occupy the ligand binding domain of the proper tyrosine kinase receptors on the target cell surface, but fail to activate cell growth.

Alternatively, antagonists to the polypeptides of the present invention may be employed which bind to the receptors to which a polypeptide of the present invention normally binds. The antagonists may be closely related proteins such that they recognize and bind to the receptor sites of the natural protein, however, they are

inactive forms of the natural protein and thereby prevent the action of VEGF-2 since receptor sites are occupied. In these ways, the action of the VEGF-2 is prevented and the antagonist/inhibitors may be used therapeutically as an anti-tumor drug by occupying the receptor sites of tumors which are recognized by VEGF-2 or by inactivating VEGF-2 itself. The antagonist/inhibitors may also be used to prevent inflammation due to the increased vascular permeability action of VEGF-2. The antagonist/inhibitors may also be used to treat solid tumor growth, diabetic retinopathy, psoriasis and rheumatoid arthritis.

The antagonist/inhibitors may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinabove described.

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

In order to facilitate understanding of the following examples, certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 mg of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 Fl of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 mg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37EC are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel, D. *et al.*, *Nucleic Acids Res.* 8:4057 (1980).

"Oligonucleotides" refer to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands, which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5 mg of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described by the method of Graham, F. and Van der Eb, A., *Virology* 52:456-457 (1973).

Examples

Example 1

Expression Pattern of VEGF-2 in Human Tissues and Breast Cancer Cell Lines

Northern blot analysis was carried out to examine the levels of expression of VEGF-2 in human tissues and breast cancer cell lines in human tissues. Total cellular RNA samples were isolated with RNazol™ B system (Biotech Laboratories, Inc.). About 10 mg of total RNA isolated from each breast tissue and cell line specified was separated on 1% agarose gel and blotted onto a nylon filter, (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989)). The labeling reaction was done according to the Stratagene Prime-It kit with 50 ng DNA fragment. The labeled DNA was purified with a Select-G-50 column from 5 Prime + 3 Prime, Inc (Boulder, CO). The filter was then hybridized with a radioactive labeled full length VEGF-2 gene at 1,000,000 cpm/ml in 0.5 M NaPO₄ and 7 % SDS overnight at 65°C. After washing

twice at room temperature and twice at 60°C with 0.5 X SSC, 0.1 % SDS, the filters were then exposed at -70°C overnight with an intensifying screen. A message of 1.6 Kd was observed in 2 breast cancer cell lines. Figure 5, lane #4 represents a very tumorigenic cell line that is estrogen independent for growth.

Also, 10 mg of total RNA from 10 human adult tissues were separated on an agarose gel and blotted onto a nylon filter. The filter was then hybridized with radioactively labeled VEGF-2 probe in 7% SDS, 0.5 M NaPO₄, pH 7.2; 1% BSA overnight at 65°C. Following washing in 0.2 X SSC at 65°C, the filter was exposed to film for 24 days at -70°C with intensifying screen. See Figure 6.

Example 2

Expression of the Truncated Form of VEGF-2 (SEQ ID NO:4) by in vitro Transcription and Translation

The VEGF-2 cDNA was transcribed and translated *in vitro* to determine the size of the translatable polypeptide encoded by the truncated form of VEGF-2 and a partial VEGF-2 cDNA. The two inserts of VEGF-2 in the pBluescript SK vector were amplified by PCR with three pairs of primers, 1) M13-reverse and forward primers; 2) M13-reverse primer and VEGF primer F4; and 3) M13-reverse primer and VEGF primer F5. The sequence of these primers are as follows.

M13-2 reverse primer: 5'-ATGCTTCCGGCTCGTATG-3' (SEQ ID NO:9)

This sequence is located upstream of the 5' end of the VEGF-2 cDNA insert in the pBluescript vector and is in an anti-sense orientation as the cDNA. A T3 promoter sequence is located between this primer and the VEGF-2 cDNA.

M13-2 forward primer: 5'GGGTTTTCCAGTCACGAC-3' (SEQ ID NO:10)

This sequence is located downstream of the 3' end of the VEGF-2 cDNA insert in the pBluescript vector and is in an anti-sense orientation as the cDNA insert.

VEGF primer F4: 5'-CCACATGGTTCAGGAAAGACA-3' (SEQ ID NO:11)

This sequence is located within the VEGF-2 cDNA in an anti-sense orientation from bp 1259-1239, which is about 169 bp away from the 3' end of the stop codon and about 266 bp before the last nucleotide of the cDNA.

PCR reaction with all three pairs of primers produce amplified products with T3 promoter sequence in front of the cDNA insert. The first and third pairs of primers produce PCR products that encode the polypeptide of VEGF-2 shown in SEQ ID NO:4. The second pair of primers produce PCR product that misses 36 amino acids coding sequence at the C-terminus of the VEGF-2 polypeptide.

Approximately 0.5 mg of PCR product from first pair of primers, 1 mg from second pair of primers, 1 mg from third pair of primers were used for *in vitro* transcription/translation. The *in vitro* transcription/translation reaction was performed in a 25 Fl of volume, using the T_NTJ Coupled Reticulocyte Lysate Systems (Promega, CAT# L4950). Specifically, the reaction contains 12.5 Fl of T_NT rabbit reticulocyte lysate 2 Fl of T_NT reaction buffer, 1 Fl of T3 polymerase, 1 Fl of 1 mM amino acid mixture (minus methionine), 4 Fl of ³⁵S-methionine (>1000 Ci/mmol, 10 mCi/ml), 1 Fl of 40 U/μl; RNasin ribonuclease inhibitor, 0.5 or 1 mg of PCR products. Nuclease-free H₂O was added to bring the volume to 25 Fl. The reaction was incubated at 30°C for 2 hours. Five microliters of the reaction product was analyzed on a 4-20% gradient SDS-PAGE gel. After fixing in 25% isopropanol and 10% acetic acid, the gel was dried and exposed to an X-ray film overnight at 70°C.

As shown in Figure 7, PCR products containing the truncated VEGF-2 cDNA (*i.e.*, as depicted in SEQ ID NO:3) and the cDNA missing 266 bp in the 3' untranslated region (3'-UTR) produced the same length of translated products, whose molecular weights are estimated to be 38-40 dk (lanes 1 and 3). The cDNA missing all the 3'UTR and missing sequence encoding the C-terminal 36 amino acids was translated into a polypeptide with an estimated molecular weight of 36-38 kd (lane 2).

Example 3 Cloning and Expression of VEGF-2 Using the Baculovirus Expression System

The DNA sequence encoding the VEGF-2 protein without 46 amino acids at the N-terminus, see ATCC No. 97149, was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

The 5' primer has the sequence TGT AAT ACG ACT CAC TAT AGG GAT CCC GCC ATG GAG GCC ACG GCT TAT GC (SEQ ID NO:12) and contains a BamH1 restriction enzyme site (in bold) and 17 nucleotide sequence complementary to the 5' sequence of VEGF-2 (nt. 150-166).

The 3' primer has the sequence GATC TCT AGA TTA GCT CAT TTG TGG TCT (SEQ ID NO:13) and contains the cleavage site for the restriction enzyme XbaI and 18 nucleotides complementary to the 3' sequence of VEGF-2, including the stop codon and 15 nt sequence before stop codon.

The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101, Inc., La Jolla, CA). The fragment was then digested with the endonuclease BamH1 and XbaI and then purified again on

a 1% agarose gel. This fragment was ligated to pAcGP67A baculovirus transfer vector (Pharmingen) at the BamHI and XbaI sites. Through this ligation, VEGF-2 cDNA was cloned in frame with the signal sequence of baculovirus gp67 gene and was located at the 3' end of the signal sequence in the vector. This is designated pAcGP67A-VEGF-2.

To clone VEGF-2 with the signal sequence of gp67 gene to the pRG1 vector for expression, VEGF-2 with the signal sequence and some upstream sequence were excised from the pAcGP67A-VEGF-2 plasmid at the Xho restriction endonuclease site located upstream of the VEGF-2 cDNA and at the XbaI restriction endonuclease site by XhoI and XbaI restriction enzyme. This fragment was separated from the rest of vector on a 1% agarose gel and was purified using "GeneClean" kit. It was designated F2.

The PRG1 vector (modification of pVL941 vector) is used for the expression of the VEGF-2 protein using the baculovirus expression system (for review see: Summers, M.D. and Smith, G.E., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555, (1987)). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by the recognition sites for the restriction endonucleases BamHI, SmaI, XbaI, BglII and Asp718. A site for restriction endonuclease XhoI is located upstream of BamHI site. The sequence between XhoI and BamHI is the same as that in pAcGp67A (static on tape) vector. The polyadenylation site of the simian virus (SV)40 is used for efficient polyadenylation. For an easy selection of recombinant virus the beta-galactosidase gene from *E. coli* is inserted in the same orientation as the polyhedrin promoter followed by the polyadenylation signal of the polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for the cell-mediated homologous recombination of cotransfected wild-type viral DNA. Many other baculovirus vectors could be used in place of pRG1 such as pAc373, pVL941 and pAcIM1 (Luckow, V.A. and Summers, M.D., *Virology* 170:31-39 (1989)).

The plasmid was digested with the restriction enzymes XhoI and XbaI and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The DNA was then isolated from a 1% agarose gel using the commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated V2.

Fragment F2 and the dephosphorylated plasmid V2 were ligated with T4 DNA ligase. *E. coli* HB101 cells were then transformed and bacteria identified that

contained the plasmid (pBac gp67-VEGF-2) with the VEGF-2 gene using the enzymes BamHI and XbaI. The sequence of the cloned fragment was confirmed by DNA sequencing.

5 5 mg of the plasmid pBac gp67-VEGF-2 was cotransfected with 1.0 mg of a commercially available linearized baculovirus ("BaculoGoldJ baculovirus DNA", Pharmingen, San Diego, CA.) using the lipofectin method (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987)).

10 1mg of BaculoGoldJ virus DNA and 5 mg of the plasmid pBac gp67-VEGF-2 were mixed in a sterile well of a microtiter plate containing 50 μ l of serum free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10 μ l Lipofectin plus 90 μ l Grace's medium were added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture was added dropwise to the Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate was rocked back and forth to mix the
15 newly added solution. The plate was then incubated for 5 hours at 27°C. After 5 hours the transfection solution was removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum was added. The plate was put back into an incubator and cultivation continued at 27°C for four days.

After four days the supernatant was collected and a plaque assay performed
20 similar as described by Summers and Smith, *supra*. As a modification an agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) was used which allows an easy isolation of blue stained plaques. (A detailed description of a "plaque assay" can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

25 Four days after the serial dilution, the virus was added to the cells, blue stained plaques were picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses was then resuspended in an Eppendorf tube containing 200 μ l of Grace's medium. The agar was removed by a brief centrifugation and the supernatant containing the recombinant baculovirus was used to infect Sf9 cells
30 seeded in 35 mm dishes. Four days later the supernatants of these culture dishes were harvested and then stored at 4°C.

Sf9 cells were grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells were infected with the recombinant baculovirus V-gp67-VEGF-2 at a multiplicity of infection (MOI) of 1. Six hours later the medium was
35 removed and replaced with SF900 II medium minus methionine and cysteine (Life Technologies Inc., Gaithersburg). 42 hours later 5 mCi of ³⁵S-methionine and 5 mCi

³⁵S cysteine (Amersham) were added. The cells were further incubated for 16 hours before they were harvested by centrifugation and the labelled proteins visualized by SDS-PAGE and autoradiography.

Protein from the medium and cytoplasm of the Sf9 cells was analyzed by SDS-PAGE under non-reducing and reducing conditions. See Figures 8A and 8B, respectively. The medium was dialyzed against 50 mM MES, pH 5.8. Precipitates were obtained after dialysis and resuspended in 100 mM NaCitrate, pH 5.0. The resuspended precipitate was analyzed again by SDS-PAGE and was stained with Coomassie Brilliant Blue. See Figure 9.

The medium supernatant was also diluted 1:10 in 50 mM MES, pH 5.8 and applied to an SP-650M column (1.0 x 6.6 cm, Toyopearl) at a flow rate of 1 ml/min. Protein was eluted with step gradients at 200, 300 and 500 mM NaCl. The VEGF-2 was obtained using the elution at 500 mM. The eluate was analyzed by SDS-PAGE in the presence or absence of reducing agent, b-mercaptoethanol and stained by Coomassie Brilliant Blue. See Figure 10.

Example 4

Expression of Recombinant VEGF-2 in COS Cells

The expression of plasmid, VEGF-2-HA is derived from a vector pcDNA1/Amp (Invitrogen) containing: 1) SV40 origin of replication, 2) ampicillin resistance gene, 3) *E. coli* replication origin, 4) CMV promoter followed by a polylinker region, an SV40 intron and polyadenylation site. A DNA fragment encoding the entire VEGF-2 precursor and a HA tag fused in frame to its 3' end was cloned into the polylinker region of the vector, therefore, the recombinant protein expression is directed under the CMV promoter. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein as previously described (Wilson *et al.*, *Cell* 37:767 (1984)). The infusion of HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is described as follows:

The DNA sequence encoding VEGF-2, ATCC No. 97149, was constructed by PCR using two primers: the 5' primer (CGC GGA TCC ATG ACT GTA CTC TAC CCA) (SEQ ID NO:14) contains a BamH1 site followed by 18 nucleotides of VEGF-2 coding sequence starting from the initiation codon; the 3' sequence (CGC TCT AGA TCA AGC GTA GTC TGG GAC GTC GTA TGG GTA CTC GAG GCT

CAT TTG TGG TCT 3') (SEQ ID NO:15) contains complementary sequences to an XbaI site, HA tag, XhoI site, and the last 15 nucleotides of the VEGF-2 coding sequence (not including the stop codon). Therefore, the PCR product contains a BamHI site, coding sequence followed by an XhoI restriction endonuclease site and HA tag fused in frame, a translation termination stop codon next to the HA tag, and an XbaI site. The PCR amplified DNA fragment and the vector, pcDNAI/Amp, were digested with BamHI and XbaI restriction enzyme and ligated. The ligation mixture was transformed into *E. coli* strain SURE (Stratagene Cloning Systems, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant VEGF-2, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Laboratory Press, (1989)). The expression of the VEGF-2-HA protein was detected by radiolabelling and immunoprecipitation method (E. Harlow and D. Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post transfection. Culture media was then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson *et al.*, *Cell* 37:767 (1984)). Both cell lysate and culture media were precipitated with an HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.

Example 5
The Effect of Partially Purified VEGF-2 Protein on the Growth of Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) were seeded at 2×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium was replaced with M199 containing 10% FBS, 8 units/ml heparin. VEGF-2 protein of SEQ ID NO. 2 minus the initial 45 amino acid residues, (VEGF) and basic FGF (bFGF) were added, at the concentration shown. On days 4 and 6, the medium was replaced. On day 8, cell number was determined with a Coulter Counter (See Figure 12).

Example 6
The Effect of Purified VEGF-2 Protein on the Growth of Vascular Endothelial Cells

5 On day 1, human umbilical vein endothelial cells (HUVEC) were seeded at $2-5 \times 10^4$ cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium was replaced with M199 containing 10% FBS, 8 units/ml heparin. Purified VEGF-2 protein of SEQ ID NO:2 minus initial 45
10 amino acid residues was added to the medium at this point. On days 4 and 6, the medium was replaced with fresh medium and supplements. On day 8, cell number was determined with a Coulter Counter (See Figure 13).

Example 7
Expression via Gene Therapy

15 Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at
20 room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin, is added. This is then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks
25 in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

30 pMV-7 (Kirschmeier, P.T. *et al.*, DNA 7:219-225 (1988) flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

35 The cDNA encoding a polypeptide of the present invention is amplified using PCR primers which correspond to the 5' and 3' end sequences respectively. The 5' primer containing an EcoRI site and the 3' primer further includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is used to transform bacteria HB101,

which are then plated onto agar-containing kanamycin for the purpose of confirming that the vector had the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells are transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as *neo* or *his*.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product.

Example 8

Expression of VEGF-2 mRNA in Human Fetal and Adult Tissues

Experimental Design

Northern blot analysis was carried out to examine the levels of expression of VEGF-2 mRNA in human fetal and adult tissues. A cDNA probe containing the entire nucleotide sequence of the VEGF-2 protein was labeled with ^{32}P using the rediprime[®] DNA labeling system (Amersham Life Science), according to the manufacturer's instructions. After labeling, the probe was purified using a CHROMA SPIN-100* column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe was then used to examine various human tissues for VEGF-2 mRNA.

A Multiple Tissue Northern (MTN) blot containing various human tissues (Fetal Kidney, Fetal Lung, Fetal Liver, Brain, Kidney, Lung, Liver, Spleen, Thymus, Bone Marrow, Testes, Placenta, and Skeletal Muscle) was obtained from

Clontech. The MTN blot was examined with the labeled probe using ExpressHyb* hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blot was exposed to film at -70°C overnight with an intensifying screen and developed according to standard procedures.

Results

Expression of VEGF-2 mRNA is abundant in vascular smooth muscle and several highly vascularized tissues. VEGF-2 is expressed at significantly higher levels in tissues associated with hematopoietic or angiogenic activities, i.e. fetal kidney, fetal lung, bone marrow, placental, spleen and lung tissue. The expression level of VEGF-2 is low in adult kidney, fetal liver, adult liver, testes; and is almost undetectable in fetal brain, and adult brain (See Figure 14).

In primary cultured cells, the expression of VEGF-2 mRNA is abundant in vascular smooth muscle cells and dermal fibroblast cells, but much lower in human umbilical vein endothelial cells (see Figure 15). This mRNA distribution pattern is very similar to that of VEGF.

Example 9

Construction of Amino terminal and carboxy terminal deletion mutants

In order to identify and analyze biologically active VEGF-2 polypeptides, a panel of deletion mutants of VEGF-2 was constructed using the expression vector pHE4a.

1. Construction of VEGF-2 T103-L215 in pHE4

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 T103-L215 (amino acids 103 to 215 in Figure 1 or SEQ ID NO:18) into the E. coli protein expression vector, pHE4, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

5' Primer (Nde I/START and 18 nt of coding sequence):

5'-GCA GCA CAT ATG ACA GAA GAG ACT ATA AAA-3' (SEQ ID NO:

19)

3' Primer (Asp718, STOP, and 15 nt of coding sequence):

5'-GCA GCA GGT ACC TCA CAG TTT AGA CAT GCA-3' (SEQ ID NO: 20).

5 The above described 5' primer (SEQ ID NO: 19), incorporates an NdeI restriction site and the above described 3' Primer (SEQ ID NO:20), incorporates an Asp718 restriction site. The 5' primer (SEQ ID NO:19) also contains an ATG sequence adjacent and in frame with the VEGF-2 coding region to allow translation of the cloned fragment in E. coli, while the 3' primer (SEQ ID NO:20) contains one stop codon (preferentially utilized in E. coli) adjacent and in frame with the VEGF-2 coding region which ensures correct translational termination in E. coli.

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419 in SEQ ID NO:18) as, for example, constructed in Example 3 as template.

15 The resulting amplicon was restriction digested with NdeI and Asp718 and subcloned into NdeI/Asp718 digested pHE4a expression vector.

2. Construction of VEGF-2 T103-R227 in pHE4

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 T103-R227 (amino acids 103 to 227 in Figure 1 or SEQ ID NO:18) into the E. coli protein expression vector, pHE4, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

25 5' Primer (Nde I/START and 18 nt of coding sequence):

5'-GCA GCA CAT ATG ACA GAA GAG ACT ATA AAA-3' (SEQ ID NO:19)

3' Primer (Asp 718, STOP, and 15 nt of coding sequence):

30 5'-GCA GCA GGT ACC TCA ACG TCT AAT AAT GGA-3' (SEQ ID NO:21)

In the case of the above described primers, an NdeI or Asp718 restriction site was incorporated into the 5' primer and 3' primer, respectively. The 5' primer (SEQ ID NO:19) also contains an ATG sequence adjacent and in frame with the VEGF-2 coding region to allow translation of the cloned fragment in E. coli, while the 3'

35

Primer (SEQ ID NO:21) contains one stop codon (preferentially utilized in *E. coli*) adjacent and in frame with the VEGF-2 coding region which ensures correct translational termination in *E. coli*.

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419 in SEQ ID NO:18) as, for example, constructed in Example 3, as template. The resulting amplicon was restriction digested with NdeI and Asp718 and subcloned into NdeI/Asp718 digested pHE4a protein expression vector.

3. Construction of VEGF-2 T103-L215 in pA2GP

In this illustrative example, the plasmid shuttle vector pA2 GP is used to insert the cloned DNA encoding the N-terminal and C-terminal deleted VEGF-2 protein (amino acids 103-215 in Figure 1 or SEQ ID NO:18), into a baculovirus to express the N-terminal and C-terminal deleted VEGF-2 protein, using a baculovirus leader and standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by the secretory signal peptide (leader) of the baculovirus gp67 protein and convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that expresses the cloned polynucleotide.

Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow *et al.*, *Virology* 170:31-39 (1989).

The cDNA sequence encoding the VEGF-2 protein without 102 amino acids at the N-terminus and without 204 amino acids at the C-terminus in Figure 1, was

amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene.

The 5' primer has the sequence 5' -GCA GCA **GGA TCC CAC AGA AGA GAC TAT AAA**- 3' (SEQ ID NO:22) containing the BamHI restriction enzyme site (in bold) followed by 1 spacer nt to stay in-frame with the vector-supplied signal peptide, and 17 nt of coding sequence bases of VEGF-2 protein. The 3' primer has the sequence 5N-GCA GCA **TCT AGA TCA CAG TTT AGA CAT GCA**-3' (SEQ ID NO:23) containing the XbaI restriction site (in bold) followed by a stop codon and 17 nucleotides complementary to the 3' coding sequence of VEGF-2.

The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101, Inc., La Jolla, CA). The fragment was then digested with the endonuclease BamHI and XbaI and then purified again on a 1% agarose gel. This fragment was ligated to pA2 GP baculovirus transfer vector (Supplier) at the BamHI and XbaI sites. Through this ligation, VEGF-2 cDNA representing the N-terminal and C-terminal deleted VEGF-2 protein (amino acids 103-215 in Figure 1 or SEQ ID NO:18) was cloned in frame with the signal sequence of baculovirus GP gene and was located at the 3' end of the signal sequence in the vector. This is designated pA2GPVEGF-2.T103-L215.

4. Construction of VEGF-2 T103-R227 in pA2GP

The cDNA sequence encoding the VEGF-2 protein without 102 amino acids at the N-terminus and without 192 amino acids at the C-terminus in Figure 1 (i.e., amino acids 103-227 of SEQ ID NO:18) was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene.

The 5'-GCA GCA **GGA TCC CAC AGA AGA GAC TAT AAA ATT TGC TGC**-3' primer has the sequence (SEQ ID NO:24) containing the BamHI restriction enzyme site (in bold) followed by 1 spacer nt to stay in-frame with the vector-supplied signal peptide, and 26 nt of coding sequence bases of VEGF-2 protein. The 3' primer has the sequence 5N-GCA GCA **TCT AGA TCA ACG TCT AAT AAT GGA ATG AAC**-3' (SEQ ID NO:25) containing the XbaI restriction site (in bold) followed by a stop codon and 21 nucleotides complementary to the 3' coding sequence of VEGF-2.

The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101, Inc., La Jolla, CA). The fragment was then digested with the endonuclease BamHI and XbaI and then purified again on a 1% agarose gel. This fragment was ligated to pA2 GP baculovirus transfer vector (Supplier) at the BamHI and XbaI sites. Through this ligation, VEGF-2 cDNA

representing the N-terminal and C-terminal deleted VEGF-2 protein (amino acids 103-227 in Figure 1 or SEQ ID NO:18) was cloned in frame with the signal sequence of baculovirus GP gene and was located at the 3' end of the signal sequence in the vector. This construct is designated pA2GPVEGF-2.T103-R227.

5. Construction of VEGF-2 in pC1

The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology*, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3N intron, the polyadenylation and termination signal of the rat preproinsulin gene.

The vector pC1 is used for the expression of VEGF-2 protein. Plasmid pC1 is a derivative of the plasmid pSV2-dhfr [ATCC Accession No. 37146]. Both plasmids contain the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., Alt, F.W., Kellems, R.M., Bertino, J.R., and Schimke, R.T., 1978, *J. Biol. Chem.* 253:1357-1370, Hamlin, J.L. and Ma, C. 1990, *Biochem. et Biophys. Acta*, 1097:107-143, Page, M.J. and Sydenham, M.A. 1991, *Biotechnology* 9:64-68). Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene it is usually co-amplified and over-expressed. It is state of the art to develop cell lines carrying more than 1,000 copies of the genes. Subsequently, when the methotrexate is withdrawn, cell lines contain the amplified gene integrated into the chromosome(s).

Plasmid pC1 contains for the expression of the gene of interest a strong promoter of the long terminal repeat (LTR) of the Rouse Sarcoma Virus (Cullen, *et al.*, *Molecular and Cellular Biology*, March 1985:438-4470) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530, 1985). Downstream of the promoter are the following single restriction enzyme cleavage sites that allow the integration of the

genes: BamHI, PvuII, and NruI. Behind these cloning sites the plasmid contains translational stop codons in all three reading frames followed by the 3'N intron and the polyadenylation site of the rat preproinsulin gene. Other high efficient promoters can also be used for the expression, e.g., the human b-actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV-I. For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well.

Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

The plasmid pC1 is digested with the restriction enzyme BamHI and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

The DNA sequence encoding VEGF-2, ATCC Accession No. 97149, was constructed by PCR using two primers corresponding to the 5' and 3' ends of the VEGF-2 gene: the 5' Primer (5'-GAT CGA TCC ATC ATG CAC TCG CTG GGC TTC TTC TCT GTG GCG TGT TCT CTG CTC G-3' (SEQ ID NO:26)) contains a Klenow-filled BamHI site and 40 nt of VEGF-2 coding sequence starting from the initiation codon; the 3' primer (5'-GCA GGG TAC GGA TCC TAG ATT AGC TCA TTT GTG GTC TTT-3' (SEQ ID NO:27)) contains a BamHI site and 16 nt of VEGF-2 coding sequence not including the stop codon.

The PCR amplified DNA fragment is isolated from a 1% agarose gel as described above and then digested with the endonuclease BamHI and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 cells are then transformed and bacteria identified that contained the plasmid pC1. The sequence and orientation of the inserted gene is confirmed by DNA sequencing. This construct is designated pC1VEGF-2.

6. Construction of pC4SigVEGF-2 T103-L215

Plasmid pC4Sig is plasmid pC4 (Accession No. 209646) containing a human IgG Fc portion as well as a protein signal sequence.

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 T103-L215 (amino acids 103 to 215 in Figure 1 or SEQ ID NO:18) into

pC4Sig, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

5' Primer (Bam HI and 26 nt of coding sequence):

5'-GCA GCA GGA TCC ACA GAA GAG ACT ATA AAA TTT GCT GC-3'
(SEQ ID NO:34)

3' Primer (Xba I, STOP, and 15 nt of coding sequence):

5'-CGT CGT TCT AGA TCA CAG TTT AGA CAT GCA TCG GCA G-3'
(SEQ ID NO:35)

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419) as, for example, constructed in Example 3, as template. The resulting amplicon was restriction digested with BamHI and XbaI and subcloned into BamHI/XbaI digested pC4Sig vector.

7. Construction of pC4SigVEGF-2 T103-R227

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 T103-L215 (amino acids 103 to 227 in Figure 1 or SEQ ID NO:18) into pC4Sig, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

5' Primer (Bam HI and 26 nt of coding sequence):

5'-GCA GCA GGA TCC ACA GAA GAG ACT ATA AAA TTT GCT GC-3'
(SEQ ID NO: 34)

3' Primer (Xba I, STOP, and 21 nt of coding sequence):

5'-GCA GCA TCT AGA TCA ACG TCT AAT AAT GGA ATG AAC-3'
(SEQ ID NO:25)

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419) as, for example, constructed in Example 3, as template. The resulting amplicon was restriction digested with BamHI and XbaI and subcloned into BamHI/XbaI digested pC4Sig vector.

8. Construction of pC4VEGF-2 M1-M263

The expression vector pC4 contains the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology*, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vector contains in addition the 3N intron, the polyadenylation and termination signal of the rat preproinsulin gene.

In this illustrative example, the cloned DNA encoding the C-terminal deleted VEGF-2 M1-M263 protein (amino acids 1-263 in Figure 1 or SEQ ID NO:18) is inserted into the plasmid vector pC4 to express the C-terminal deleted VEGF-2 protein.

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 M1-M263 into the expression vector, pC4, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

5' Primer 5'-GAC TGG ATC CGC CAC CAT GCA CTC GCT GGG CTT CTT CTC-3' (SEQ ID NO:28)

3' Primer 5'-GAC TGG TAC CTT ATC ACA TAA AAT CTT CCT GAG CC-3' (SEQ ID NO:29)

In the case of the above described 5' primer, an BamHI restriction site was incorporated, while in the case of the 3' primer, an Asp718 restriction site was incorporated. The 5' primer also contains 6 nt, 20 nt of VEGF-2 coding sequence, and an ATG sequence adjacent and in frame with the VEGF-2 coding region to allow translation of the cloned fragment in *E. coli*, while the 3' primer contains 2 nt, 20 nt of VEGF-2 coding sequence, and one stop codon (preferentially utilized in *E. coli*) adjacent and in frame with the VEGF-2 coding region which ensures correct translational termination in *E. coli*.

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419) as constructed, for example, in Example 3 as template. The resulting amplicon was restriction digested with BamHI and Asp718 and subcloned into

BamH1/Asp718 digested pC4 protein expression vector. This construct is designated pC4VEGF-2 M1-M263.

9. Construction of pC4VEGF-2 M1-D311

5 In this illustrative example, the cloned DNA encoding the C-terminal deleted VEGF-2 M1-D311 protein (amino acids 1-311 in Figure 1 or SEQ ID NO:18) is inserted into the plasmid vector pC4 to express the C-terminal deleted VEGF-2 protein.

10 To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 M1-D311 into the expression vector, pC4, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

15 5' Primer 5'-GAC TGG ATC CGC CAC CAT GCA CTC GCT GGG CTT CTT CTC-3' (SEQ ID NO:30)

3' Primer 5'-GAC TGG TAC CTT ATC AGT CTA GTT CTT TGT GGG G-3' (SEQ ID NO:31)

20 In the case of the above described 5' primer, an BamH1 restriction site was incorporated, while in the case of the 3' primer, an Asp718 restriction site was incorporated. The 5' primer also contains 6 nt, 20 nt of VEGF-2 coding sequence, and an ATG sequence adjacent and in frame with the VEGF-2 coding region to allow translation of the cloned fragment in E. coli, while the 3' primer contains 2 nt, 20 nt
25 of VEGF-2 coding sequence, and one stop codon (preferentially utilized in E. coli) adjacent and in frame with the VEGF-2 coding region which ensures correct translational termination in E. coli.

30 The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419) as constructed, for example, in Example 3 as template. The resulting amplicon was restriction digested with BamH1 and Asp718 and subcloned into BamH1/Asp718 digested pC4 protein expression vector.

10. Construction of pC4VEGF-2 M1-Q367

In this illustrative example, the cloned DNA encoding the C-terminal deleted VEGF-2 M1-D311 protein (amino acids 1-311 in SEQ ID NO:18) is inserted into the plasmid vector pC4 to express the C-terminal deleted VEGF-2 protein.

To permit Polymerase Chain Reaction directed amplification and sub-cloning of VEGF-2 M1-D311 into the expression vector, pC4, two oligonucleotide primers complementary to the desired region of VEGF-2 were synthesized with the following base sequence:

5' Primer 5'-GAC TGG ATC CGC CAC CAT GCA CTC GCT GGG CTT CTT CTC-3' (SEQ ID NO:32)

3' Primer 5'-GAC TGG TAC CTC ATT ACT GTG GAC TTT CTG TAC ATT C-3' (SEQ ID NO:33)

In the case of the above described 5' primer, an BamHI restriction site was incorporated, while in the case of the 3' primer, an Asp718 restriction site was incorporated. The 5' primer also contains 6 nt, 20 nt of VEGF-2 coding sequence, and an ATG sequence adjacent and in frame with the VEGF-2 coding region to allow translation of the cloned fragment in E. coli, while the 3' primer contains 2 nt, 20 nt of VEGF-2 coding sequence, and one stop codon (preferentially utilized in E. coli) adjacent and in frame with the VEGF-2 coding region which ensures correct translational termination in E. coli.

The Polymerase Chain Reaction was performed using standard conditions well known to those skilled in the art and the nucleotide sequence for the mature VEGF-2 (aa 24-419) as constructed, for example, in Example 3 as template. The resulting amplicon was restriction digested with BamHI and Asp718 and subcloned into BamHI/Asp718 digested pC4 protein expression vector. This construct is designated pC4VEGF-2 M1-Q367.

Example 10

Transient Expression of VEGF-2 Protein in COS-7 Cells

Experimental Design

Expression of the VEGF-2-HA fusion protein from the construct made in Example 4, for example, was detected by radiolabeling and immunoprecipitation,

using methods described in, for example Harlow and colleagues (Antibodies: A Laboratory Manual, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988)). To this end, two days after transfection, the cells were labeled by incubation in media containing 35S-cysteine for 8 hours. The cells and the media were collected, and the cells were washed and then lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson and colleagues (supra). Proteins were precipitated from the cell lysate and from the culture media using an HA-specific monoclonal antibody. The precipitated proteins then were analyzed by SDS-PAGE and autoradiography.

Results

As shown in Figure 16, cells transfected with pcDNA1 VEGF-2HA secreted a 56 kd and a 30 kd protein. The 56 kd protein, but not the 30 kd protein, could also be detected in the cell lysate but is not detected in controls. This suggests the 30 kd protein is likely to result from cleavage of the 56 kd protein. Since the HA-tag is on the C-terminus of VEGF-2, the 30 kd protein must represent the C-terminal portion of the cleaved protein, whereas the N-terminal portion of the cleaved protein would not be detected by immunoprecipitation. These data indicate that VEGF-2 protein expressed in mammalian cells is secreted and processed.

Example 11

Stimulatory effect of VEGF-2 on proliferation of vascular endothelial cells

Experimental Design

Expression of VEGF-2 is abundant in highly vascularized tissues. Therefore the role of VEGF-2 in regulating proliferation of several types of endothelial cells was examined.

Endothelial cell proliferation assay

For evaluation of mitogenic activity of growth factors, the colorimetric MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was performed (CellTiter 96 AQ, Promega). Cells were seeded in a 96-well plate (5,000 cells/well) in 0.1 mL serum-supplemented medium and allowed to attach

overnight. After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF₁₆₅ or VEGF-2 in 0.5% FBS) with or without Heparin (8 U/ml) were added to wells for 48 hours. 20 mg of MTS/PMS mixture (1:0.05) were added per well and allowed to incubate for 1 hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader. Background absorbance from control wells (some media, no cells) was subtracted, and seven wells were performed in parallel for each condition. See, Leak *et al. In Vitro Cell. Dev. Biol.* 30A:512-518 (1994)

Results

VEGF-2 stimulated proliferation of human umbilical vein endothelial cells (HUVEC) and dermal microvascular endothelial cells slightly (Figures 17 and 18). The stimulatory effect of VEGF-2 is more pronounced on proliferation of endometrial and microvascular endothelial cells (Figure 19). Endometrial endothelial cells (HEEC) demonstrated the greatest response to VEGF-2 (96% of the effect of VEGF on microvascular endothelial cells). The response of microvascular endothelial cells (HMEC) to VEGF-2 was 73% compared to VEGF. The response of HUVEC and BAEC (bovine aortic endothelial cells) to VEGF-2 was substantially lower at 10% and 7%, respectively. The activity of VEGF-2 protein has varied between different purification runs with the stimulatory effect of certain batches on HUVEC proliferation being significantly higher than that of other batches.

Example 12

Inhibition of PDGF-induced vascular smooth muscle cell proliferation

VEGF-2 expression is high in vascular smooth muscle cells. Smooth muscle is an important therapeutic target for vascular diseases, such as restenosis. To evaluate the potential effects of VEGF-2 on smooth muscle cells, the effect of VEGF-2 on human aortic smooth muscle cell (HAoSMC) proliferation was examined.

Experimental Design

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation. Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6 mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse anti-BrdUrd antibody at 4 °C for 2 h after exposing to

denaturing solution and then with the streptavidin-peroxidase and diaminobenzidine. After counterstaining with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-positive cells to the total cell number. In addition, the simultaneous detection of the BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for individual cells by the concomitant use of bright field illumination and dark field-UV fluorescent illumination. See, Hayashida *et al.*, *J. Biol. Chem.* 6;271(36):21985-21992 (1996).

Results

VEGF-2 has an inhibitory effect on proliferation of vascular smooth muscle cells induced by PDGF, but not by Fetal Bovine Serum (FBS) (Figure 20).

Example 13

Stimulation of endothelial cell migration

Endothelial cell migration is an important step involved in angiogenesis.

Experimental Design

This example will be used to explore the possibility that VEGF-2 may stimulate lymphatic endothelial cell migration. Currently, there are no published reports of such a model. However, we will be adapting a model of vascular endothelial cell migration for use with lymphatic endothelial cells essentially as follows:

Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD; Falk, W., Goodwin, R. H. J., and Leonard, E. J. "A 48 well micro chemotaxis assembly for rapid and accurate measurement of leukocyte migration." *J. Immunological Methods* 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8 μ m (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours at room temperature and dried under sterile air. Test substances are diluted to appropriate concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25 μ l of the final dilution is placed in the lower chamber of the modified Boyden apparatus. Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber, 2.5×10^5 cells suspended in 50 μ l M199 containing 1% FBS are seeded in the upper compartment. The apparatus

is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO₂ to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.

Results

In an assay examining HUVEC migration using a 43-well microchemotaxis chamber, VEGF-2 was able to stimulate migration of HUVEC (Figure 21).

Example 14


Stimulation of nitric oxide production by endothelial cells

Nitric oxide released by the vascular endothelium is believed to be a mediator of vascular endothelium relaxation. VEGF-1 has been demonstrated to induce nitric oxide production by endothelial cells in response to VEGF-1. As a result, VEGF-2 activity can be assayed by determining nitric oxide production by endothelial cells in response to VEGF-2.

Experimental Design

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of VEGF-1 and VEGF-2. Nitric oxide in the medium is determined by use of the Griess reagent to measure total nitrite after reduction of nitric oxide-derived nitrate by nitrate reductase. The effect of VEGF-2 on nitric oxide release was examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer was measured with a NO-specific polarographic electrode connected to a NO meter (Iso-NO, World Precision Instruments Inc.) (1049). Calibration of the NO elements was performed according to the following equation:



The standard calibration curve was obtained by adding graded concentrations of KNO₂ (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing KI and H₂SO₄. The specificity of the Iso NO electrode to NO was previously determined by measurement of NO from authentic NO gas (1050). The

culture medium was removed and HUVECs were washed twice with Dulbecco's phosphate buffered saline. The cells were then bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates were kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C. The NO sensor probe was inserted vertically into the wells, keeping the tip of the electrode 2 mm under the surface of the solution, before addition of the different conditions. S-nitroso acetyl penicillamin (SNAP) was used as a positive control. The amount of released NO was expressed as picomoles per 1×10^6 endothelial cells. All values reported were means of four to six measurements in each group (number of cell culture wells). See, Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

Results

VEGF-2 was capable of stimulating nitric oxide release on HUVEC (Figure 22) to a higher level than VEGF. This suggested that VEGF-2 may modify vascular permeability and vessel dilation.

Example 15

Effect of VEGF-2 on cord formation in angiogenesis

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to form capillary-like structures (hollow structures) when cultured *in vitro*.

Experimental Design

CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 μ l/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 μ g Cell Applications' Chord Formation Medium containing control buffer or HGS protein (0.1 to 100 ng/ml) and the cells are cultured for an additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

Commercial (R&D) VEGF (50 ng/ml) is used as a positive control. b-esteradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

5 Results

It has been observed that VEGF-2 inhibits cord formation similar to IFN α which also stimulates endothelial cell proliferation (Figure 23). This inhibitory effect may be a secondary effect of endothelial proliferation which is mutually exclusive with the cord formation process.

10

Example 16

Angiogenic effect on chick chorioallantoic membrane

Chick chorioallantoic membrane (CAM) is a well-established system to
15 examine angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of VEGF-2 to stimulate angiogenesis in CAM was examined.

Experimental Design

Embryos

20 Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese quail (*Coturnix coturnix*) were incubated at 37.8°C and 80% humidity. Differentiated CAM of 16-day-old chick and 13-day-old quail embryos was studied with the following methods.

25 CAM Assay

On Day 4 of development, a window was made into the egg shell of chick eggs. The embryos were checked for normal development and the eggs sealed with cellotape. They were further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) were cut into disks of about 5 mm in diameter. Sterile and salt-free
30 growth factors were dissolved in distilled water and about 3.3 mg/ 5 ml was pipetted on the disks. After air-drying, the inverted disks were applied on CAM. After 3 days, the specimens were fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They were photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described
35 above. Controls were performed with carrier disks alone.

Results

This data demonstrates that VEGF-2 can stimulate angiogenesis in the CAM assay nine-fold compared to the untreated control. However, this stimulation is only 45% of the level of VEGF stimulation (Figure 24).

5

Example 17

Angiogenesis assay using a Matrigel implant in mouse

Experimental Design

10 In order to establish an *in vivo* model for angiogenesis to test protein activities, mice and rats have been implanted subcutaneously with methylcellulose disks containing either 20 mg of BSA (negative control) and 1 mg of bFGF and 0.5 mg of VEGF-1 (positive control).

15 It appeared as though the BSA disks contained little vascularization, while the positive control disks showed signs of vessel formation. At day 9, one mouse showed a clear response to the bFGF.

Results

20 Both VEGF proteins appeared to enhance Matrigel cellularity by a factor of approximately 2 by visual estimation.

An additional 30 mice were implanted with disks containing BSA, bFGF, and varying amounts of VEGF-1, VEGF-2-B8, and VEGF-2-C4. Each mouse received two identical disks, rather than one control and one experimental disk.

25 Samples of all the disks recovered were immunostained with Von Willebrand's factor to detect for the presence of endothelial cells in the disks, and flk-1 and flt-4 to distinguish between vascular and lymphatic endothelial cells. However, definitive histochemical analysis of neovascularization and lymphangiogenesis could not be determined.

30

Example 18

Rescue of Ischemia in Rabbit Lower Limb Model

Experimental Design

35 To study the *in vivo* effects of VEGF-2 on ischemia, a rabbit hindlimb ischemia model was created by surgical removal of one femoral arteries as described previously (Takeshita, S. *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). The excision

of the femoral artery results in retrograde propagation of thrombus and occlusion of the external iliac artery. Consequently, blood flow to the ischemic limb is dependent upon collateral vessels originating from the internal iliac artery (Takeshita, S. *et al. Am J. Pathol* 147:1649-1660 (1995)). An interval of 10 days was allowed for post-operative recovery of rabbits and development of endogenous collateral vessels. At 10 day post-operatively (day 0), after performing a baseline angiogram, the internal iliac artery of the ischemic limb was transfected with 500 mg naked VEGF-2 expression plasmid by arterial gene transfer technology using a hydrogel-coated balloon catheter as described (Riessen, R. *et al. Hum Gene Ther.* 4:749-758 (1993); Leclerc, G. *et al. J. Clin. Invest.* 90: 936-944 (1992)). When VEGF-2 was used in the treatment, a single bolus of 500 mg VEGF-2 protein or control was delivered into the internal iliac artery of the ischemic limb over a period of 1 min. through an infusion catheter. On day 30, various parameters were measured in these rabbits.

15 Results

Both VEGF-2 protein (Figure 25, top panels) and naked expression plasmid (Figure 25, middle panels) were able to restore the following parameters in the ischemic limb. Restoration of blood flow, angiographic score seem to be slightly more by administration of 500 mg plasmid compared with by 500 mg protein (Figure 25, bottom panels) The extent of the restoration is comparable with that by VEGF in separate experiments (data not shown). A vessel dilator was not able to achieve the same effect, suggesting that the blood flow restoration is not simply due to a vascular dilation effect.

25 a. BP ratio (Figure 25a)

The blood pressure ratio of systolic pressure of the ischemic limb to that of normal limb.

2. Blood Flow and Flow Reserve (Figure 25b)

30 Resting FL: the blood flow during un-dilated condition

Max FL: the blood flow during fully dilated condition (also an indirect measure of the blood vessel amount)

Flow Reserve is reflected by the ratio of max FL: resting FL.

35 3 Angiographic Score (Figure 25c)

This is measured by the angiogram of collateral vessels. A score was determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh.

5 4. Capillary density (Figure 25d)

The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

10 As discussed, VEGF-2 is processed to an N-terminal and a C-terminal fragment which are co-purified. The N-terminal fragment contains the intact putative functional domain and may be responsible for the biologic activity.

Example 19

Effect of VEGF-2 on Vasodilation

15

As described above, VEGF-2 can stimulate NO release, a mediator of vascular endothelium dilation. Since dilation of vascular endothelium is important in reducing blood pressure, the ability of VEGF-2 to affect the blood pressure in spontaneously hypertensive rats (SHR) was examined. VEGF-2 caused a dose-dependent decrease
20 in diastolic blood pressure (Figures 26a and b). There was a steady decline in diastolic blood pressure with increasing doses of VEGF-2 which attained statistical significance when a dose of 300mg/kg was administered. The changes observed at this dose were not different than those seen with acetylcholine (0.5mg/kg). Decreased mean arterial pressure (MAP) was observed as well (Figure 26c and d). VEGF-2
25 (300 mg/kg) and acetylcholine reduced the MAP of these SHR animals to normal levels.

Additionally, increasing doses (0, 10, 30, 100, 300, and 900 mg/kg) of the B8, C5, and C4 preps of VEGF-2 were administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM.
30 Statistical analysis was performed with a paired t-test and statistical significance was defined as $p < 0.05$ vs. the response to buffer alone.

Studies with VEGF-2 (C5 prep) revealed that although it significantly decreased the blood pressure, the magnitude of the response was not as great as that seen with VEGF-2 (B8 prep) even when used at a dose of 900 mg/kg.

35 Studies with VEGF-2 (C4 preparation) revealed that this CHO expressed protein preparation yielded similar results to that seen with C5 (i.e. statistically

significant but of far less magnitude than seen with the B8 preparation) (see Figures 26A-D).

As a control and since the C4 and C5 batches of VEGF-2 yielded minor, but statistically significant, changes in blood pressure, experiments were performed
5 experiments with another CHO-expressed protein, M-CIF. Administration of M-CIF at doses ranging from 10-900 mg/kg produced no significant changes in diastolic blood pressure. A minor statistically significant reduction in mean arterial blood pressure was observed at doses of 100 and 900 mg/kg but no dose response was noted. These results suggest that the reductions in blood pressure observed with the
10 C4 and C5 batches of VEGF-2 were specific, i.e. VEGF-2 related.

Example 20

Rat Ischemic Skin Flap Model

15 **Experimental Design**

The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. VEGF-2 expression, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- 20
- a) Ischemic skin
 - b) Ischemic skin wounds
 - c) Normal wounds

The experimental protocol includes:

- 25
- a) Raising a 3x4 cm, single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).
 - b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).
 - c) Topical treatment with VEGF-2 of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.
 - 30 d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

Example 21***Peripheral Arterial Disease Model***

Angiogenic therapy using VEGF-2 has been developed as a novel therapeutic strategy to obtain restoration of blood flow around the ischemia in case of peripheral arterial diseases.

Experimental Design

The experimental protocol includes:

- a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.
- b) VEGF-2 protein, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3 weeks.
- c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of VEGF-2 expression and histology. Biopsy is also performed on the other side of normal muscle of the contralateral hindlimb.

Example 22***Ischemic Myocardial Disease Model***

VEGF-2 is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after coronary artery occlusion. Alteration of VEGF-2 expression is investigated in situ.

Experimental Design

The experimental protocol includes:

- a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.
- b) VEGF-2 protein, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4 weeks.
- c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

Example 23**Rat Corneal Wound Healing Model**

This animal model shows the effect of VEGF-2 on neovascularization.

5~

Experimental Design

The experimental protocol includes:

- a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.
- b) Inserting a spatula below the lip of the incision facing the outer corner of the eye.
- 10 c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).
- d) Positioning a pellet, containing 50mg - 500mg VEGF-2, within the pocket.
- e) VEGF-2 treatment can also be applied topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

15

Example 24**Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models****Experimental Design**

20 The experimental protocol includes:

1. Diabetic db+/db+ mouse model.

To demonstrate that VEGF-2 accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

30 The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy,

myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

Animals

Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates were used in this study (Jackson Laboratories). The animals were purchased at 6 weeks of age and were 8 weeks old at the beginning of the study. Animals were individually housed and received food and water ad libitum. All manipulations were performed using aseptic techniques. The experiments were conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Surgical Wounding

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two-day intervals thereafter. Wound closure is determined by daily

measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

- 5 VEGF-2 is administered using at a range different doses of VEGF-2, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

- 10 Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Experimental Design

- 15 Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) were evaluated: 1) Vehicle placebo control, 2) VEGF-2.

Measurement of Wound Area and Closure

- 20 Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 was 64mm², the corresponding size of the dermal punch. Calculations were made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

25

Histology

- 30 Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with KGF-2. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.
- 35

Immunohistochemistry

Re-epithelialization

Tissue sections are stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used
5 as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

Cell Proliferation Marker

10 Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer served as a positive tissue control and human brain tissue is used as a negative tissue control. Each specimen included a section with omission of the primary antibody and substitution with non-immune mouse IgG.
15 Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Statistical Analysis

20 Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

B. Steroid Impaired Rat Model

The inhibition of wound healing by steroids has been well documented in
25 various *in vitro* and *in vivo* systems (Wahl, S.M. Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl, S.M.*et al.*, *J. Immunol.* 115: 476-481 (1975); Werb, Z. *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert, R.H., *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis
30 (Beck, L.S. *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes, B.F. *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes, B.F., *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, S. M., "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic
35 and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish

phenomenon in rats (Beck, L.S. *et al.*, *Growth Factors*. 5: 295-304 (1991); Haynes, B.F., *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, S. M., "Glucocorticoids and wound healing", *In: Antiinflammatory Steroid Action: Basic and Clinical Aspects*, Academic Press, New York, pp. 280-302 (1989); Pierce, G.F. *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that VEGF-2 can accelerate the healing process, the effects of multiple topical applications of VEGF-2 on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Animals

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and were 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Surgical Wounding

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8 for Figure. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if

granulation tissue was no longer visible and the wound is covered by a continuous epithelium.

VEGF-2 is administered using at a range different doses of VEGF-2, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Experimental Design

Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) were evaluated: 1) Untreated group 2) Vehicle placebo control 3) VEGF-2 treated groups.

Measurement of Wound Area and Closure

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 was 64mm², the corresponding size of the dermal punch. Calculations were made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Histology

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining was performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin was improved by treatment with VEGF-2. A calibrated lens micrometer was used by a blinded observer to determine the distance of the wound gap.

Statistical Analysis

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

Example 25
Specific Peptide Fragments to Generate
VEGF-2 Monoclonal Antibodies

Four specific peptides (designated SP-40, SP-41, SP-42 and SP-43) have been generated. These will be used to generate monoclonal antibodies to analyze VEGF-2 processing. The peptides are shown below:

1. "SP-40": MTVLYPEYWKMY (amino acids 70-81 in SEQ ID NO:18)
2. "SP-41": KSIDNEWRKTSMPREV (amino acids 120-136(note C->S mutation at position 131) in SEQ ID NO: 18)
3. "SP-42": MSKLDVYRQVHSIIRR (amino acids 212-227 in SEQ ID NO: 18)
4. "SP-43": MFSSDAGDDSTDGFHDI (amino acids 263-279 in SEQ ID NO: 18)

Example 26
Lymphadema Animal Model

The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of VEGF-2 in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

Experimental Procedure

Prior to beginning surgery, blood sample was drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs were shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements were made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of

1% Evan's Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats
5 are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated or suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located.
10 The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care was taken to control any mild bleeding resulting from this procedure. After lymphatics were occluded, the skin flaps are sealed by using liquid skin
15 (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals were checked daily through the optimal edematous peak, which
20 typically occurred by day 5-7. The plateau edematous peak was then observed. To evaluate the intensity of the lymphedema, we measured the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days. The effect plasma proteins have on lymphedema and determined if protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous
25 limbs are evaluated at 2 places. Analysis is performed in a blind manner.

Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged.
30 Readings are taken from both control and edematous limbs.

Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), both legs are
35 shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco

edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

5 Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca²⁺ comparison.

10 Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs were amputated using a quillitine, then both experimental and control legs were cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint was disarticulated and the foot was weighed.

15 Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle was observed under fluorescent microscopy for lymphatics. Other immuno/histological methods are currently being evaluated.

Example 27

20 Method of Treatment Using Gene Therapy for Production of VEGF-2 Polypeptide - In Vivo

25 Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) comprising VEGF-2 operably linked to a promoter into an animal to increase the expression of VEGF-2. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO 90/11092, WO 98/11779; U.S. Patent Nos. 5693622, 5705151, 5580859; Tabata H. *et al.* (1997) *Cardiovasc. Res.* 35(3):470-479, Chao, J *et al.* (1997) *Pharmacol. Res.* 35(6):517-522, Wolff, J.A. (1997) *Neuromuscul. Disord.* 7(5):314-318, Schwartz, B. *et al.* (1996) *Gene Ther.* 3(5):405-411, Tsurumi, Y. *et al.* (1996) *Circulation* 94(12):3281-3290 (incorporated herein by reference).

30 The VEGF-2 polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The VEGF-2 polynucleotide constructs may also be delivered directly into arteries.

The VEGF-2 polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the VEGF-2 polynucleotide may also be delivered in liposome formulations (such as those taught in Felgner P.L. *et al.* (1995) *Ann. NY Acad. Sci.* 772:126-139 and Abdallah B. *et al.* (1995) *Biol. Cell* 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The VEGF-2 vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The VEGF-2 construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. Preferably, they are delivered by direct injection into the artery.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the

tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues, or directly into arteries. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked VEGF-2 constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

10 The dose response effects of injected VEGF-2 polynucleotide construct in arteries *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for VEGF-2 is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The arteries of rabbits are
15 then injected with various amounts of the template DNA.

Hindlimb ischemia in rabbits is surgically induced, as described in Example 18. Immediately following this, five different sites in the adductor (2 sites), medial large (2 sites), and semimembranous muscles (1 site) are injected directly with plasmid DNA encoding VEGF-2 using a 3ml syringe and 2-gauge needle advanced
20 through a small skin incision. The skin is then closed using 4.0 nylon.

The ability to rescue hindlimb ischemia is determined by measuring the number of capillaries in light microscopic sections taken from the treated hindlimbs, compared to ischemic hindlimbs from untreated rabbits, measurement of calf blood pressure, and intra-arterial Doppler guidewire measurement of flow velocity
25 (Takeshita *et al.*, *J. Clin. Invest.* 93:662-670 (1994)). The results of the above experimentation in rabbits can be used to extrapolate proper dosages and other treatment parameters in humans and other animals using VEGF-2 polynucleotide naked DNA.

Example 28

30 Method of Treatment Using Gene Therapy - Ex Vivo

One method of gene therapy transplants fibroblasts, which are capable of expressing VEGF-2 polypeptides, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet
35 surface of a tissue culture flask, approximately ten pieces are placed in each flask.

The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding VEGF-2 can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector contains properly inserted VEGF-2.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the VEGF-2 gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the VEGF-2 gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently

infected, the fibroblasts are analyzed to determine whether VEGF-2 protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads

5

Example. 29

10

Method of Treatment Using Gene Therapy Homologous Recombination

Another method of gene therapy according to the present invention involves operably associating the endogenous VEGF-2 sequence with a promoter via
15 homologous recombination as described, for example, in U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature* 342:435-438 (1989). This method involves the activation of a gene
20 which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous VEGF-2, flanking the promoter. The targeting sequence will be sufficiently near the
25 5' end of VEGF-2 so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the
30 amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added
35 together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous VEGF-2 sequence. This results in the expression of VEGF-2 in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na_2HPO_4 , 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

Plasmid DNA is prepared according to standard techniques. To construct a plasmid for targeting to the VEGF-2 locus, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two VEGF-2 non-coding sequences are amplified via PCR: one VEGF-2 non-coding sequence (VEGF-2 fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other VEGF-2 non-coding sequence (VEGF-2 fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and VEGF-2 fragments are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; VEGF-2 fragment 1 - XbaI; VEGF-2 fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120 $\mu\text{g/ml}$. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is

performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 μ F and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37EC. The following day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product.

Example 30

VEGF-2 Transgenic Animals

The VEGF-2 polypeptides can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i.e., polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science

259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred.

Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis

of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

5 Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher
10 levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the
15 transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of VEGF-2 polypeptides, studying conditions and/or disorders associated with aberrant
20 VEGF-2 expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.-

Example 31

VEGF-2 Knock-Out Animals

25 Endogenous VEGF-2 gene expression can also be reduced by inactivating or "knocking out" the VEGF-2 gene and/or its promoter using targeted homologous recombination. (*E.g.*, see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of
30 which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express
35 polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene

of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas & 5 Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to 10 express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (*e.g.*, knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (*i.e.*, animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (*e.g.*, lymphocytes), adipocytes, 15 muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, *e.g.*, by transduction (using viral vectors, and preferably vectors that integrate the 20 transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the VEGF-2 polypeptides. The engineered 25 cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, *e.g.*, in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, *e.g.*, genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or 30 vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson; U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the 35 development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an

exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Knock-out animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of VEGF-2 polypeptides, studying conditions and/or disorders associated with aberrant VEGF-2 expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

145

Applicant's or agent's file reference number	PF112PCT3	International application No. PCT/US 99/05021
-------------------------------------------------	-----------	--------------------------------------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>3</u> , line <u>24</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 12, 1995	Accession Number 97149
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application Authorized officer / Jeff Meadows Paraegal Specialist IAPD-PCT Operations (703) 306-3745	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on: Authorized officer

Applicant's or agent's file reference number	PF112PCT3	146 International Patent Application	PCT/US 9 / 05-021
----------------------------------------------	-----------	-----------------------------------------	-------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>3</u> , line <u>25</u>					
B. IDENTIFICATION OF DEPOSIT					
Further deposits are identified on an additional sheet <input type="checkbox"/>					
Name of depositary institution American Type Culture Collection					
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America					
Date of deposit March 4, 1994	Accession Number 75698				
C. ADDITIONAL INDICATIONS (leave blank if not applicable)					
This information is continued on an additional sheet <input type="checkbox"/>					
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)					
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)					
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")					
<table border="1"> <tr> <td>For receiving Office use only</td> <td>For International Bureau use only</td> </tr> <tr> <td> <input checked="" type="checkbox"/> This sheet was received with the international application Lydel Meadows Legal Specialist Authorized officer IAPD-PCT Operations (703) 305-3745 </td> <td> <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer </td> </tr> </table>		For receiving Office use only	For International Bureau use only	<input checked="" type="checkbox"/> This sheet was received with the international application Lydel Meadows Legal Specialist Authorized officer IAPD-PCT Operations (703) 305-3745	<input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer
For receiving Office use only	For International Bureau use only				
<input checked="" type="checkbox"/> This sheet was received with the international application Lydel Meadows Legal Specialist Authorized officer IAPD-PCT Operations (703) 305-3745	<input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer				

What Is Claimed Is:

1. An isolated polynucleotide comprising a nucleic acid sequence at least 95% identical to a polynucleotide which encodes a polypeptide selected from the group consisting of:

(a) a N-terminal deletion fragment described by the general formula m-396 of SEQ ID NO:2;

(b) a C-terminal deletion fragment described by the general formula -23-n of SEQ ID NO:2;

(c) a N-terminal and C-terminal deletion fragment described by the general formula m-n of SEQ ID NO:2; and

(d) a C-terminal deletion fragment described by the general formula +9-n of SEQ ID NO:2.

2. The isolated polynucleotide of claim 1, wherein said polypeptide comprises amino acid residues S-205 to S-396 of SEQ ID NO:2 and/or amino acid residues F-9 to R-203 of SEQ ID NO:2.

3. A composition comprising the isolated polynucleotide of claim 1.

4. The isolated polynucleotide of claim 1, wherein the polynucleotide encodes a biologically active fragment of VEGF-2.

5. The isolated polynucleotide of claim 1, wherein the polynucleotide encodes a polypeptide which binds an antibody for VEGF-2.

6. The polynucleotide of claim 1 further comprising a polynucleotide which encodes a heterologous polypeptide.

7. A vector comprising the polynucleotide of claim 1.

8. The vector of claim 7, wherein said polynucleotide is operatively associated with a heterologous regulatory sequence.

9. A host cell comprising the vector of claim 7 or the polynucleotide of claim 1.

10. A method for producing a VEGF-2 polypeptide, comprising:
 - (a) culturing the host cell of claim 9 under conditions suitable to produce the polypeptide; and
 - (b) recovering the polypeptide from the cell culture.
11. The polypeptide produced by the method of claim 10.
12. An isolated polypeptide comprising polypeptide at least 95% identical to an amino acid sequence selected from the group consisting of:
 - (a) a N-terminal deletion fragment described by the general formula m-396 of SEQ ID NO:2;
 - (b) a C-terminal deletion fragment described by the general formula -23-n of SEQ ID NO:2;
 - (c) a N-terminal and C-terminal deletion fragment described by the general formula m-n of SEQ ID NO:2; and
 - (d) a C-terminal deletion fragment described by the general formula +9-n of SEQ ID NO:2.
13. The isolated polypeptide of claim 12, wherein said polypeptide comprises amino acid residues S-205 to S-396 of SEQ ID NO:2 and/or amino acid residues F-9 to R-203 of SEQ ID NO:2.
14. A composition comprising the isolated polypeptide of claim 1.
15. A composition comprising a first polypeptide fragment comprising amino acids residues S-205 to S-396 of SEQ ID NO:2 and a second polypeptide fragment comprising amino acid residues F-9 to R-203 of SEQ ID NO:2.
16. The isolated polypeptide of claim 12, wherein the polypeptide is a biologically active fragment.
17. The isolated polypeptide of claim 12, wherein the polypeptide is antigenic.

18. The isolated polypeptide of claim 12, further comprising a heterologous polypeptide.

19. An antibody to the polypeptide of claim 12.

20. A compound which activates the polypeptide of claim 12.

21. A compound which inhibits the polypeptide of claim 12.

22. A method for preventing, treating, or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of the polypeptide of claims 12-13, the composition of claims 3, 14-15, or of the polynucleotide of claim 1-2.

23. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of a secreted protein comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of claim 1;

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

24. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of a secreted protein comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 12 in a biological sample;

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

25. A method for identifying binding partner to the polypeptide of claim 10 comprising:

(a) contacting the polypeptide of claim 12 with a binding partner; and

(b) determining whether the binding partner effects an activity of the polypeptide.

26. A method of identifying an activity in a biological assay, wherein the method comprises:

- (a) expressing the polypeptide of claim 12 in a cell;
- (b) isolating the supernatant;
- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

FIG. 1A

1 GTCCCTCCACCATGCACTCGCTGGGCTTCTTCTGTGGGTGTTCTCTGCTCGCCGCTG 60
 CAGGAAGGTGTACGTGAGCGACCCGAAGAAGAGACACCCGACAAAGAGACGAGCGCGGAC
 M H S L G F F S V A C S L L A A A -
 C
 61 CGCTGCTCCCGGTCCTCGGAGGGCCCGCCGCGCCGCCCTTCGAGTCCGGACTCG 120
 GCGACGAGGCCCCAGGAGCGCTCCGCGGGCGGGCGGGGGAAGCTCAGGCCCTGAGC
 L L P G P R E A P A A A A F E S G L D -
 C
 121 ACCTCTCGGACGGGAGCCCGACGGCGGGGAGGCCACGGCTTATGCAAGCAAAGATCTGG 180
 TGGAGAGCCTGCGCCTCGGGCTGCGCCCGCTCCGGTGCCGAATACGTTCTGTTCTAGACC
 L S D A E P D A G E A T A Y A S K D L E -
 C
 181 AGGAGCAGTTACGGTCTGTGTCCAGTGTAGATGAACATCATGACTGTACTCTACCCAGAAT 240
 TCCTCGTCAATGCCAGACACAGGTCACTACTTGTAGTACTGACATGAGATGGGTCTTA
 E Q L R S V S S V D E L M T V L Y P E Y -
 C
 241 ATTGGAAATGTACAAGTGTGAGTAAAGAAAGGAGGCTGGCAACATAACAGAGAACAGG 300
 TAACCTTTTACATGTTACAGTCGATTCTCTTCTCCGACCGTGTATGTCTCTTGTCC
 W K M Y K C Q L R K G G W Q H N R E Q A -
 C
 CCAACCTCAACTCAAGGACAGAGAGACTATAAAATTCTGTCAGCACATTATAATACAG

MATCH WITH FIG. 1B

MATCH WITH FIG. 1A

200

U

075

—

085

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

050

1

000

1

000

1

MATCH WITH FIG. 1C

MATCH WITH FIG. 1B

MATCH WITH FIG. 1D

MATCH WITH FIG. 1C

C	V	C	K	N	K	L	F	P	S	Q	C	G	A	N	R	E	F	D	E	N	-
	ACACATGCCAGTGTATGTAAGAACCTGCCCCAGAAATCAACCCCTAAATCCTGGAA																				
	1021																				
	TGTTACGGTCACACATACATTTCTTGGACGGGCTTTAGTTGGGATTTAGGACCTT																				
	T	C	Q	C	V	C	K	R	T	C	P	R	N	Q	P	L	N	P	G	K	-
C	AATGTGCCTGTGAATGTACAGAAAGTCCACAGAAATGCTTGTAAAGGAAAGAGTTCC																				
	1140																				
	TTACACGGACACTTACATGTCCTTTCAGGTGCTTTACGAACAATTTTCTTCTTCAAGG																				
	C	A	C	E	C	T	E	S	P	Q	K	C	L	L	K	G	K	K	F	H	-
C	ACCACCAAACATGCAGCTGTTACAGACGGCCATGTACGAACCGCCAGAAAGGCTTGTGAGC																				
	1200																				
	TGGTGGTTGTACGTCGACAATGTCTGCCGGTACATGCTTGGCGGCTTCCGAACACTCG																				
	H	Q	T	C	S	C	Y	R	R	P	C	T	N	R	Q	K	A	C	E	P	-
C	CAGGATTTTCATATAGTGAAGAGTGTGTCGTTGTGTCCTTCATATTTGGCAAAGACCAC																				
	1260																				
	GTCCTAAAGTATATCACTTCTTCACACAGCAACACAGGGAAGTATAACCGTTTCTGTTG																				
	G	F	S	Y	S	E	E	V	C	R	C	V	P	S	Y	W	Q	R	P	Q	-
C	AAATGAGCTAAGATTGTACTGTTTTCAGTTTCATCGATTTTCTATATTATGGAACACTGTGT																				

MATCH WITH FIG. 1E

FIG. 1E

MATCH WITH FIG. 1D

1261 -----+-----+-----+-----+-----+-----+-----+ 1320
 TTTACTCGATTCTAACATGACAAAGGTCAAGTAGCTAAAGATAATACCTTTTGACACA
 C M S *
 1321 TGCCACAGTAGAACTGTGTGAACAGAGAGACCCTTGTGGGTCCATGCTAACAAAGACA 1380
 -----+-----+-----+-----+-----+-----+-----+
 ACGGTGTCATCTTGACAGACACTGTGCTCTCTCTGGGAACACCCAGGTACGATTGTTCTGT
 1381 AAAGTCTGTCTTTCTGAAACCATGTGGATAAATTACAGAAATGGACTGGAGCTCATCTG 1440
 -----+-----+-----+-----+-----+-----+-----+
 TTTCAGACAGAAAGGACTTGGTACACCTATTGAATGTCTTTACCTGACCTCGAGTAGAC
 CAAAAGGCCCTCTTGTAAGACTGCTTTCTGCCAATGACCAAAACAGCCAAGATTTTCCTC
 1441 -----+-----+-----+-----+-----+-----+-----+ 1500
 GTTTTCCGGAGAACATTTCTGACCAAAAGACGGTTACTGGTTTGTCTGGTTCTAAAAGGAG
 TTGTGATTTCTTTAAAGAATGACTATATAATTTATTTCCCACTAAAAATATTGTTCTGC
 1501 -----+-----+-----+-----+-----+-----+-----+ 1560
 AACACTAAAGAAATTTTCTTACTGATATATAATAAAGGTGATTTTATAACAAGACG
 ATTCATTTTATAGCAACAATAATGGTAAACTCCTGATCAATATTTTATATCAT
 1561 -----+-----+-----+-----+-----+-----+-----+ 1620
 TAAGTAAATAATCGTTGTTGTTAACCATTTTGAGTGACACTAGTTATAAAAATATAGTA
 GCAAAATATGTTTAAATAATAATGAAAATTGTATTATATAAAAAA
 1621 -----+-----+-----+-----+-----+-----+-----+ 1674
 CGTTTATACAAATTTTATTATTACCTTTTAAACATAAATAATTTT
 C

```
1  CGAGGCCACGGCTTATGCAAGCAAGATCTGGAGGAGCAGTTACGGTCTGTGTCCAGTGT  
-----+-----+-----+-----+-----+-----+-----+  
71  AGATGAACTCATGACTGTACTCTACCCAGAATATTGAAATGTACAAGTGCAGCTAAG  
-----+-----+-----+-----+-----+-----+-----+  
      M T V L Y P E Y W K M Y K C Q L R  
-----+-----+-----+-----+-----+-----+-----+  
121 GAAAGGAGGCTGGCAACATAACAGAGAACAGGCCAACCTCAACTCAAGCAGAGAGAC  
-----+-----+-----+-----+-----+-----+-----+  
      K G G W Q H N R E Q A N L N S R T E E T  
-----+-----+-----+-----+-----+-----+-----+  
181 TATAAAATTGCTGCAGCACATTATAATACAGAGATCTTGAAAAGTATTGATAATCAGTC  
-----+-----+-----+-----+-----+-----+-----+  
      I K F A A A H Y N T E I L K S I D N E W  
-----+-----+-----+-----+-----+-----+-----+  
241 GAGAAAGACTCAATGCATGCCACGGAGGTGTGTATAGATGTGGGGAAGGAGTTTGGAGT  
-----+-----+-----+-----+-----+-----+-----+  
      R K T Q C M P R E V C I D V G K E F G V  
-----+-----+-----+-----+-----+-----+-----+  
301 CGCGACAAACACCTTCTTTAAACCTCCATGTGTGTCGCTACAGATGTGGGGTTCCTG  
-----+-----+-----+-----+-----+-----+-----+  
      A T N T F F K P P C V S V Y R C G G C C
```

FIG. 2A

361 CAATAGTGGGGCTGCAGTGCATGAACACCAGCAGGCTACCTCAGCAAGACGTTATT
-----+-----+-----+-----+-----+
N S E G L Q C M N T S T S Y L S K T L F
421 TGAATTAAGTGCCTCTCTCTCAAGGCCCAACCAGTAACAATCAGTTTGGCCAATCA
-----+-----+-----+-----+-----+
E I T V P L S Q G P K P V T I S F A N H
481 CACTTCCTGCCGATGCATGTCTAACTGGATGTTTACAGACAAGTTCATTCCATTATTAG
-----+-----+-----+-----+-----+
T S C R C M S K L D V Y R Q V H S I I R
541 ACGTTCCTGCCAGCAACACTACCACAGTGTCTCAGGCAGCGAACAAGACCTGCCCCACCAA
-----+-----+-----+-----+-----+
R S L P A T L P Q C Q A A N K T C P T N
601 TTACATGTGGAATAATCACATCTGCAGATGCCCTGGCTCAGGAAGATTTATGTTTCCTC
-----+-----+-----+-----+-----+
Y M W N N H I C R C L A Q E D F M F S S
661 GGATGCTGGAGATGACTCAACAGATGGATTCCATGACATCTGTGGACCAACAAGGAGCT
-----+-----+-----+-----+-----+
D A G D D S T D G F H D I C G P N K E L

FIG.2B

721 GGATGAAGAGACCTGTCAGTGTCTGTCAGAGCGGGGCTTCGGCCTGCCAGCTGTGGACC
-----+-----+-----+-----+-----+
D E E T C Q C V C R A G L R P A S C G P
781 CCACAAAGAACTAGACAGAACTCATGCCAGTGTCTGTGTAACAACTCTTCCCCAG
-----+-----+-----+-----+-----+
H K E L D R N S C Q C V C K N K L F P S
841 CCAATGTGGGCCAACCGAGAAATTGATGAAACACATGCCAGTGTGTATGTAAGAAGAAC
-----+-----+-----+-----+-----+
Q C G A N R E F D E N T C Q C V C K R T
901 CTGCCCCAGAAATCAACCCCTAAATCCTGGAAATGTGCCCTGTGAATGTACAGAAAGTCC
-----+-----+-----+-----+-----+
C P R N Q P L N P G K C A C E C T E S P
961 ACAGAAATGCTTGTAAAGGAAGAAGTTCCACCACCAACATGCCAGCTGTTACAGACG
-----+-----+-----+-----+-----+
Q K C L L K G K K F H H Q T C S C Y R R
1021 GCCATGTACGAACCGCCAGAGGCTTGTGAGCCAGGATTTTCATATAGTGAAGAAGTGTG
-----+-----+-----+-----+-----+
P C T N R Q K A C E P G F S Y S E E V C

FIG.2C

1081 TCGTTGTGCCCTTCATATTGGCAAGACCACAAATGAGCTAAGATTGTACTGTTTCCCA
-----+-----+-----+-----+-----+-----+
R C V P S Y W Q R P Q M S
1141 GTTCATCGATTTTCTATTATGGAACACTGTGTGGCCACAGTAGAACTGCTGTGAACAGA
-----+-----+-----+-----+-----+-----+
1201 GAGACCCCTGTGGGTCCATGCTAACAAAGACAAAGTCTGTCTTTCCTGAACCATGTGGA
-----+-----+-----+-----+-----+-----+
1261 TAACTTTACAGAAATGGACTGGAGCTCATCTGCCAAAAGGCCCTCTTGTAAAGACTGGTTTT
-----+-----+-----+-----+-----+-----+
1321 CTGCCAATGACCAACAGCCAAGATTTCCTCTTGTGATTTCCTTAAAGAAATGACTATA
-----+-----+-----+-----+-----+-----+
1381 TAATTTATTTCCACTAAAATAATGTGTTCTGCATTCTATTTTATAGCAACAATTGGT
-----+-----+-----+-----+-----+-----+
1441 AAAACTCACTGTGATCAATATTTTATATCATGCAAAATATGTTTAAATAAATGAAAA
-----+-----+-----+-----+-----+-----+
1501 TTGTATTATATAAAAAAATAAAAAA
-----+-----+-----+-----+-----+-----+

FIG. 2D

1	50
pdgfa	.MRTLACLLL LGCGLAHLV AEEAEIPREV IERLARSQIH SIRDLORLLE
pdgfb	MNRCWA.LFL SLCCYLRLVS AEGDPIPEEL YEMLSDEHSIR SFDDLORLLE
VegfMNFLL SWVHWSLALL LY.....
Vegf2MTV LYPEYWKMYK CQ.....
51	100
pdgfa	IDSVGSEDSL DTSLRAHGVE ATKHVPEKRP LPIRRKRSI.EEAVP
pdgfb	GDP.GEEDGA ELDLNMTRSH SGGELES... .LARGRRSLG SLTIAEPAMI
Vegf	APMAE.....GGGQ NHHEVVKFMD .VYQR.....
Vegf2	REQANLNSRT EETIKFAAAH YNTEILKSID NEWRK.....
101	150
pdgfa	AVCKTRTVIY EIPRSQVDPT SANFLIWPPC VEVKRCTGCC NTSSVKCQPS
pdgfb	AECKTRTEVF EISRRLLDRT NANFLVWPPC VEVQRCSGCC NNRNVQCRPT
Vegf	SYCHPIETLV DIFQEYPDEI ..EYIFKPSC VPLMRCGGCC NDEGLECVPT
Vegf2	TQCMPEVCI DVGKEFGVAT ..NTFFKPPC VSVYRCGGCC NSEGLQCMNT
151	200
pdgfa	RVHRSVKVA KVEYVRKKPK LKEVQVRLEE HLECAC..... AT.....
pdgfb	QVQLRPVQVR KIEIVRKKPI FKKAIVTLED HLACKC..... ETVAARPV
Vegf	EESNITMQIM RIK.PH..QG QHIGEMSEFLQ HNKCECRPKK DRARQEKKS
Vegf2	STSYLSKTLF EIT.VPLSQG PKPVTISFAN HTSCRCMSKL DVYRQVHSII

FIG. 3A


201	250
PdgfaTSLNPD YREEDTDVR.
Pdgfb	RSPGGSQEQR AKTPQTRVTI RTVRVRPPK GKRRKFKXTH DKTALKETLG
Vegf	RGK.....GKGQKRRK KSRKSWSVY VGARCCIMPW SLPQPHP...
Vegf2	RRSLPATLPQ CQAANKTCPT NYMWNHICR CLAQEDFMFS SDAGDDSTDG
251	300
Pdgfa
Pdgfb	A.....
VegfCGP... ..CSE RRKHLEFVQDP QTCKCCKNT
Vegf2	FHDICGNKE LDEETCQCVC RAGLRPASC GPHKEL...DR NSCQCVCCKNK
301	350
Pdgfa
Pdgfb
Vegf	..DSRCKARQ LELNERTC RC DKPRR
Vegf2	LFPSQCGANR .EFDENTCQC VCKRTCPRNQ PLNPGKACE CTESPQKCLL
351	398
Pdgfa
Pdgfb
Vegf
Vegf2	KGKKFHHQTC SCYRRPCTNR QKACEPGFSY SEEVCRVPS YWQRPQMS

FIG. 3B

PERCENTAGE (%) OF AMINO ACID IDENTITIES BETWEEN EACH PAIR OF GENES IS SHOWN IN THE FOLLOWING TABLE				
	PDGF α	PDGF β	VEGF	VEGF2
PDGF α				
PDGF β	48.0			
VEGF	20.7	22.7		
VEGF2	23.5	22.4	30.0	


FIG. 4

Expression of VEGF2 mRNA in Human Breast Tumor Cells



1. normal breast tissue
2. breast tumor tissue
- 3-9. breast tumor cell lines.


FIG. 5



Expression of VEGF2 mRNA in human adult tissues.

FIG. 6

FIG. 7



Lane 1: 14-C and rainbow M.W. marker
Lane 2: FGF control
Lane 3: VEGF2 (M13-reverse & forward primers)
Lane 4: VEGF2 (M13-reverse & VEGF-F4 primers)
Lane 5: VEGF2 (M13-reverse & VEGF-F5 primers)




FIG. 8B

FIG. 8A

FIG. 10





FIG. 9



Lane 1: Molecular weight marker
Lane 2: Precipitates containing VEGF2.

FIG. 11




FIG. 12








FIG. 13



**Figure 14**

**Figure 15**

**Figure 16**







Figure 17

**Figure 18**

**Figure 19**

**Figure 20**







Figure 21

HUVEC - NO-Release**Figure 22**

**Figure 23**

**Figure 24**






Figure 25A




Figure 25B






Figure 25C









Figure 25D

**Figure 26A**

**Figure 26B**

**Figure 26C**


**Figure 26D**



The effect of increasing doses of VEGF-2 on the mean arterial pressure (MAP) of SHR rats

Figure 26E

The effect of increasing doses of VEGF-2 on the mean arterial pressure (MAP) of SHR rats




Increasing doses of VEGF-2 (HG00403-B8, HG00404-C5, and HG00404-C4) were administered to 13-14 week old SHR (spontaneously hypertensive rats) and the data were expressed as the mean \pm SEM. Statistical analysis was performed with a paired t-test and significance defined as $p < 0.05$ vs. the response to buffer alone. The response to B8 was significant at a 300 µg/kg dose. Administration of C5 yielded significant responses at doses greater than or equal to 100 µg/kg. The response to C4 was significant when 10, 100, 300, and 900 µg/kg were given.

The effect of VEGF-2 on the diastolic blood pressure of SHR rats

Figure 26F

The effect of VEGF-2 on the diastolic blood pressure of SHR rats



Increasing doses of VEGF-2 (HG00403-B8, HG00404-C5, HG00404-C4) were administered to 13-14 week old SHR (spontaneously hypertensive rats) and the data are expressed as the mean \pm SEM. Statistical analysis was performed with a paired t-test and statistical significance was defined as $p < 0.05$ vs. the response to buffer alone. The response to B8 was significant only at the 300 μ g/kg dose and when given acetylcholine. The responses to C4 and C5, while much less dramatic, were statistically significant at all dose levels.

Figure 26G




Figure 27




Figure 28

Operator 1
-35
1 AAGCTTAAAAAC TGCAAAATAGTTTGACTTTGTCAGTGGTGAATAAGCAAT

-10
Operator 2
50 TAAGATGTACCCAAATTGTTTTCACACATAA

94 A GAG GAG A A A T T A C A T A T G S/D

-1-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Human Genome Sciences, Inc.
- (ii) TITLE OF INVENTION: VASCULAR ENDOTHELIAL GROWTH FACTOR 2
- (iii) NUMBER OF SEQUENCES: 35
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: HUMAN GENOME SCIENCES, INC.
 - (B) STREET: 9410 KEY WEST AVENUE
 - (C) CITY: ROCKVILLE
 - (D) STATE: MARYLAND
 - (E) COUNTRY: USA
 - (F) ZIP: 20850
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: TO BE ASSIGNED
 - (B) FILING DATE: HERewith
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 09/042,105
 - (B) FILING DATE: 13-MAR-1998
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 09/107,997
 - (B) FILING DATE: 30-JUN-1998
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: MICHELE M. WALES
 - (B) REGISTRATION NUMBER: 43,975
 - (C) REFERENCE/DOCKET NUMBER: PF112PCT3
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (301) 309-8504
 - (B) TELEFAX: (301) 309-8439

-2-

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1674 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 12..80

(ix) FEATURE:

- (A) NAME/KEY: mat_peptide
- (B) LOCATION: 81..1268

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 12..1268

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTCCTTCCAC C ATG CAC TCG CTG GGC TTC TTC TCT GTG GCG TGT TCT CTG	50
Met His Ser Leu Gly Phe Phe Ser Val Ala Cys Ser Leu	
-23 -20 -15	
CTC GCC GCT GCG CTG CTC CCG GGT CCT CGC GAG GCG CCC GCC GCC GCC	98
Leu Ala Ala Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala	
-10 -5 1 5	
GCC GCC TTC GAG TCC GGA CTC GAC CTC TCG GAC GCG GAG CCC GAC GCG	146
Ala Ala Phe Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala	
10 15 20	
GGC GAG GCC ACG GCT TAT GCA AGC AAA GAT CTG GAG GAG CAG TTA CGG	194
Gly Glu Ala Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg	
25 30 35	
TCT GTG TCC AGT GTA GAT GAA CTC ATG ACT GTA CTC TAC CCA GAA TAT	242
Ser Val Ser Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr	
40 45 50	
TGG AAA ATG TAC AAG TGT CAG CTA AGG AAA GGA GGC TGG CAA CAT AAC	290
Trp Lys Met Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn	
55 60 65 70	
AGA GAA CAG GCC AAC CTC AAC TCA AGG ACA GAA GAG ACT ATA AAA TTT	338
Arg Glu Gln Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe	
75 80 85	

-3-

GCT GCA GCA CAT TAT AAT ACA GAG ATC TTG AAA AGT ATT GAT AAT GAG Ala Ala Ala His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu 90 95 100	386
TGG AGA AAG ACT CAA TGC ATG CCA CGG GAG GTG TGT ATA GAT GTG GGG Trp Arg Lys Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly 105 110 115	434
AAG GAG TTT GGA GTC GCG ACA AAC ACC TTC TTT AAA CCT CCA TGT GTG Lys Glu Phe Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val 120 125 130	482
TCC GTC TAC AGA TGT GGG GGT TGC TGC AAT AGT GAG GGG CTG CAG TGC Ser Val Tyr Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys 135 140 145 150	530
ATG AAC ACC AGC ACG AGC TAC CTC AGC AAG ACG TTA TTT GAA ATT ACA Met Asn Thr Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr 155 160 165	578
GTG CCT CTC TCT CAA GGC CCC AAA CCA GTA ACA ATC AGT TTT GCC AAT Val Pro Leu Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn 170 175 180	626
CAC ACT TCC TGC CGA TGC ATG TCT AAA CTG GAT GTT TAC AGA CAA GTT His Thr Ser Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val 185 190 195	674
CAT TCC ATT ATT AGA CGT TCC CTG CCA GCA ACA CTA CCA CAG TGT CAG His Ser Ile Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln 200 205 210	722
GCA GCG AAC AAG ACC TGC CCC ACC AAT TAC ATG TGG AAT AAT CAC ATC Ala Ala Asn Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile 215 220 225 230	770
TGC AGA TGC CTG GCT CAG GAA GAT TTT ATG TTT TCC TCG GAT GCT GGA Cys Arg Cys Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly 235 240 245	818
GAT GAC TCA ACA GAT GGA TTC CAT GAC ATC TGT GGA CCA AAC AAG GAG Asp Asp Ser Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu 250 255 260	866
CTG GAT GAA GAG ACC TGT CAG TGT GTC TGC AGA GCG GGG CTT CGG CCT Leu Asp Glu Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro 265 270 275	914
GCC AGC TGT GGA CCC CAC AAA GAA CTA GAC AGA AAC TCA TGC CAG TGT Ala Ser Cys Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys 280 285 290	962
GTC TGT AAA AAC AAA CTC TTC CCC AGC CAA TGT GGG GCC AAC CGA GAA Val Cys Lys Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu 295 300 305 310	1010

-4-

TTT GAT GAA AAC ACA TGC CAG TGT GTA TGT AAA AGA ACC TGC CCC AGA	1058
Phe Asp Glu Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg	
315 320 325	
AAT CAA CCC CTA AAT CCT GGA AAA TGT GCC TGT GAA TGT ACA GAA AGT	1106
Asn Gln Pro Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser	
330 335 340	
CCA CAG AAA TGC TTG TTA AAA GGA AAG AAG TTC CAC CAC CAA ACA TGC	1154
Pro Gln Lys Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys	
345 350 355	
AGC TGT TAC AGA CGG CCA TGT ACG AAC CGC CAG AAG GCT TGT GAG CCA	1202
Ser Cys Tyr Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro	
360 365 370	
GGA TTT TCA TAT AGT GAA GAA GTG TGT CGT TGT GTC CCT TCA TAT TGG	1250
Gly Phe Ser Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp	
375 380 385 390	
CAA AGA CCA CAA ATG AGC TAAGATTGTA CTGTTTTCCA GTTCATCGAT	1298
Gln Arg Pro Gln Met Ser	
395	
TTTCTATTAT GGAAACTGT GTTGCCACAG TAGAACTGTC TGTGAACAGA GAGACCCTTG	1358
TGGGTCCATG CTAACAAAGA CAAAAGTCTG TCTTTCCTGA ACCATGTGGA TAACTTTACA	1418
GAAATGGACT GGAGCTCATC TGCAAAAGGC CTCTTGTAAG GACTGGTTTT CTGCCAATGA	1478
CCAAACAGCC AAGATTTTCC TCTTGTGATT TCTTTAAAAG AATGACTATA TAATTTATTT	1538
CCACTAAAAA TATTGTTTCT GCATTCATTT TTATAGCAAC AACAATTGGT AAAACTCACT	1598
GTGATCAATA TTTTATATC ATGCAAAATA TGTTTAAAAT AAAATGAAAA TTGTATTTAT	1658
AAAAAAAAAA AAAAAA	1674

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

-5-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met His Ser Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
 -23 -20 -15 -10

Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Phe
 -5 1 5

Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
 10 15 20 25

Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 30 35 40

Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
 45 50 55

Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
 60 65 70

Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
 75 80 85

His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
 90 95 100 105

Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe
 110 115 120

Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr
 125 130 135

Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr
 140 145 150

Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
 155 160 165

Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
 170 175 180 185

Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
 190 195 200

Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
 205 210 215

Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
 220 225 230

Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
 235 240 245

Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
 250 255 260 265

-6-

Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
 270 275 280
 Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys
 285 290 295
 Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu
 300 305 310
 Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro
 315 320 325
 Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys
 330 335 340 345
 Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr
 350 355 360
 Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser
 365 370 375
 Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro
 380 385 390
 Gln Met Ser
 395

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1526 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
- (B) LOCATION: 71..142

(ix) FEATURE:

- (A) NAME/KEY: mat_peptide
- (B) LOCATION: 143..1120

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 71..1120

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGAGGCCACG GCTTATGCAA GCAAAGATCT GGAGGAGCAG TTACGGTCTG TGTCCAGTGT

60

-7-

AGATGAACTC ATG ACT GTA CTC TAC CCA GAA TAT TGG AAA ATG TAC AAG Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys-Met Tyr Lys -24 -20 -15	109
TGT CAG CTA AGG AAA GGA GGC TGG CAA CAT AAC AGA GAA CAG GCC AAC Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln Ala Asn -10 -5 1 5	157
CTC AAC TCA AGG ACA GAA GAG ACT ATA AAA TTT GCT GCA GCA CAT TAT Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr 10 15 20	205
AAT ACA GAG ATC TTG AAA AGT ATT GAT AAT GAG TGG AGA AAG ACT CAA Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln 25 30 35	253
TGC ATG CCA CGG GAG GTG TGT ATA GAT GTG GGG AAG GAG TTT GGA GTC Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val 40 45 50	301
GCG ACA AAC ACC TTC TTT AAA CCT CCA TGT GTG TCC GTC TAC AGA TGT Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys 55 60 65	349
GGG GGT TGC TGC AAT AGT GAG GGG CTG CAG TGC ATG AAC ACC AGC ACG Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr 70 75 80 85	397
AGC TAC CTC AGC AAG ACG TTA TTT GAA ATT ACA GTG CCT CTC TCT CAA Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln 90 95 100	445
GGC CCC AAA CCA GTA ACA ATC AGT TTT GCC AAT CAC ACT TCC TGC CGA Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg 105 110 115	493
TGC ATG TCT AAA CTG GAT GTT TAC AGA CAA GTT CAT TCC ATT ATT AGA Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg 120 125 130	541
CGT TCC CTG CCA GCA ACA CTA CCA CAG TGT CAG GCA GCG AAC AAG ACC Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr 135 140 145	589
TGC CCC ACC AAT TAC ATG TGG AAT AAT CAC ATC TGC AGA TGC CTG GCT Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys Leu Ala 150 155 160 165	637
CAG GAA GAT TTT ATG TTT TCC TCG GAT GCT GGA GAT GAC TCA ACA GAT Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser Thr Asp 170 175 180	685
GGA TTC CAT GAC ATC TGT GGA CCA AAC AAG GAG CTG GAT GAA GAG ACC Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr 185 190 195	733

-8-

TGT CAG TGT GTC TGC AGA GCG GGG CTT CGG CCT GCC AGC TGT GGA CCC	781
Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys Gly Pro	
200 205 210	
CAC AAA GAA CTA GAC AGA AAC TCA TGC CAG TGT GTC TGT AAA AAC AAA	829
His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys Asn Lys	
215 220 225	
CTC TTC CCC AGC CAA TGT GGG GCC AAC CGA GAA TTT GAT GAA AAC ACA	877
Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr	
230 235 240 245	
TGC CAG TGT GTA TGT AAA AGA ACC TGC CCC AGA AAT CAA CCC CTA AAT	925
Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn	
250 255 260	
CCT GGA AAA TGT GCC TGT GAA TGT ACA GAA AGT CCA CAG AAA TGC TTG	973
Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys Cys Leu	
265 270 275	
TTA AAA GGA AAG AAG TTC CAC CAC CAA ACA TGC AGC TGT TAC AGA CGG	1021
Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg	
280 285 290	
CCA TGT ACG AAC CGC CAG AAG GCT TGT GAG CCA GGA TTT TCA TAT AGT	1069
Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser Tyr Ser	
295 300 305	
GAA GAA GTG TGT CGT TGT GTC CCT TCA TAT TGG CAA AGA CCA CAA ATG	1117
Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro Gln Met	
310 315 320 325	
AGC TAAGATTGTA CTGTTTTCCA GTTCATCGAT TTTCTATTAT GGAAACTGT	1170
Ser	
GTGCCACAG TAGAACTGTC TGTGAACAGA GAGACCCTTG TGGGTCCATG CTAACAAAGA	1230
CAAAAGTCTG TCTTTCCTGA ACCATGTGGA TAACTTTACA GAAATGGACT GGAGCTCATC	1290
TGCAAAAGGC CTCTTGTAAG GACTGGTTTT CTGCCAATGA CCAAACAGCC AAGATTTTCC	1350
TCTTGTGATT TCTTTAAAG AATGACTATA TAATTTATTT CCACTAAAAA TATTGTTTCT	1410
GCATTCATTT TTATAGCAAC AACAATTGGT AAAACTCACT GTGATCAATA TTTTATATC	1470
ATGCAAAATA TGTTTAAAT AAAATGAAAA TTGTATTTAT AAAAAAAAAA AAAAAA	1526

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

-9-

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met Tyr Lys Cys Gln Leu
-24          -20          -15          -10

Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln Ala Asn Leu Asn Ser
      -5          1          5

Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr Asn Thr Glu
      10          15          20

Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln Cys Met Pro
      25          30          35          40

Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val Ala Thr Asn
      45          50          55

Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys Gly Gly Cys
      60          65          70

Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr Ser Tyr Leu
      75          80          85

Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln Gly Pro Lys
      90          95          100

Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met Ser
      105          110          115          120

Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg Arg Ser Leu
      125          130          135

Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr Cys Pro Thr
      140          145          150

Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys Leu Ala Gln Glu Asp
      155          160          165

Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser Thr Asp Gly Phe His
      170          175          180

Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr Cys Gln Cys
      185          190          195          200

Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys Gly Pro His Lys Glu
      205          210          215

Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys Asn Lys Leu Phe Pro
      220          225          230

Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr Cys Gln Cys
      235          240          245

```

-10-

Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn Pro Gly Lys
 250 255 260

Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys Cys Leu Leu Lys Gly
 265 270 275 280

Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg Pro Cys Thr
 285 290 295

Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser Tyr Ser Glu Val
 300 305 310

Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro Gln Met Ser
 315 320 325

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 196 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Arg Thr Leu Ala Cys Leu Leu Leu Leu Gly Cys Gly Tyr Leu Ala
 1 5 10 15

His Val Leu Ala Glu Glu Ala Glu Ile Pro Arg Glu Val Ile Glu Arg
 20 25 30

Leu Ala Arg Ser Gln Ile His Ser Ile Arg Asp Leu Gln Arg Leu Leu
 35 40 45

Glu Ile Asp Ser Val Gly Ser Glu Asp Ser Leu Asp Thr Ser Leu Arg
 50 55 60

Ala His Gly Val His Ala Thr Lys His Val Pro Glu Lys Arg Pro Leu
 65 70 75 80

Pro Ile Arg Arg Lys Arg Ser Ile Glu Glu Ala Val Pro Ala Val Cys
 85 90 95

Lys Thr Arg Thr Val Ile Tyr Glu Ile Pro Arg Ser Gln Val Asp Pro
 100 105 110

Thr Ser Ala Asn Phe Leu Ile Trp Pro Pro Cys Val Glu Val Lys Arg
 115 120 125

Cys Thr Gly Cys Cys Asn Thr Ser Ser Val Lys Cys Gln Pro Ser Arg
 130 135 140

-11-

Val His His Arg Ser Val Lys Val Ala Lys Val Glu Tyr Val Arg Lys
 145 150 155 160

Lys Pro Lys Leu Lys Glu Val Gln Val Arg Leu Glu Glu His Leu Glu
 165 170 175

Cys Ala Cys Ala Thr Thr Ser Leu Asn Pro Asp Tyr Arg Glu Glu Asp
 180 185 190

Thr Asp Val Arg
 195

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 241 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asn Arg Cys Trp Ala Leu Phe Leu Ser Leu Cys Cys Tyr Leu Arg
 1 5 10 15

Leu Val Ser Ala Glu Gly Asp Pro Ile Pro Glu Glu Leu Tyr Glu Met
 20 25 30

Leu Ser Asp His Ser Ile Arg Ser Phe Asp Asp Leu Gln Arg Leu Leu
 35 40 45

His Gly Asp Pro Gly Glu Glu Asp Gly Ala Glu Leu Asp Leu Asn Met
 50 55 60

Thr Arg Ser His Ser Gly Gly Glu Leu Glu Ser Leu Ala Arg Gly Arg
 65 70 75 80

Arg Ser Leu Gly Ser Leu Thr Ile Ala Glu Pro Ala Met Ile Ala Glu
 85 90 95

Cys Lys Thr Arg Thr Glu Val Phe Glu Ile Ser Arg Arg Leu Ile Asp
 100 105 110

Arg Thr Asn Ala Asn Phe Leu Val Trp Pro Pro Cys Val Glu Val Gln
 115 120 125

Arg Cys Ser Gly Cys Cys Asn Asn Arg Asn Val Gln Cys Arg Pro Thr
 130 135 140

Gln Val Gln Leu Arg Pro Val Gln Val Arg Lys Ile Glu Ile Val Arg
 145 150 155 160

-12-

Lys Lys Pro Ile Phe Lys Lys Ala Thr Val Thr Leu Glu Asp His Leu
 165 170 175
 Ala Cys Lys Cys Glu Thr Val Ala Ala Arg Pro Val Thr Arg Ser
 180 185 190
 Pro Gly Gly Ser Gln Glu Gln Arg Ala Lys Thr Pro Gln Thr Arg Val
 195 200 205
 Thr Ile Arg Thr Val Arg Val Arg Arg Pro Pro Lys Gly Lys His Arg
 210 215 220
 Lys Phe Lys His Thr His Asp Lys Thr Ala Leu Lys Glu Thr Leu Gly
 225 230 235 240
 Ala

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 232 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Asn Phe Leu Leu Ser Trp Val His Trp Ser Leu Ala Leu Leu Leu
 1 5 10 15
 Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Met Ala Glu Gly
 20 25 30
 Gly Gly Gln Asn His His Glu Val Val Lys Phe Met Asp Val Tyr Gln
 35 40 45
 Arg Ser Tyr Cys His Pro Ile Glu Thr Leu Val Asp Ile Phe Gln Glu
 50 55 60
 Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser Cys Val Pro Leu
 65 70 75 80
 Met Arg Cys Gly Gly Cys Cys Asn Asp Glu Gly Leu Glu Cys Val Pro
 85 90 95
 Thr Glu Glu Ser Asn Ile Thr Met Gln Ile Met Arg Ile Lys Pro His
 100 105 110
 Gln Gly Gln His Ile Gly Glu Met Ser Phe Leu Gln His Asn Lys Cys
 115 120 125

-13-

Glu Cys Arg Pro Lys Lys Asp Arg Ala Arg Gln Glu Lys Lys Ser Val
 130 135 140

Arg Gly Lys Gly Lys Gly Gln Lys Arg Lys Arg Lys Lys Ser Arg Tyr
 145 150 155 160

Lys Ser Trp Ser Val Tyr Val Gly Ala Arg Cys Cys Leu Met Pro Trp
 165 170 175

Ser Leu Pro Gly Pro His Pro Cys Gly Pro Cys Ser Glu Arg Arg Lys
 180 185 190

His Leu Phe Val Gln Asp Pro Gln Thr Cys Lys Cys Ser Cys Lys Asn
 195 200 205

Thr Asp Ser Arg Cys Lys Ala Arg Gln Leu Glu Leu Asn Glu Arg Thr
 210 215 220

Cys Arg Cys Asp Lys Pro Arg Arg
 225 230

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Pro Xaa Cys Val Xaa Xaa Xaa Arg Cys Xaa Gly Cys Cys Asn
 1 5 10

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGCTTCCGG CTCGTATG

18

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:

-14-

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGGTTTTCCC AGTCACGAC

18

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCACATGGTT CAGGAAAGAC A

21

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 50 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TGTAATACGA CTCACTATAG GGATCCCGCC ATGGAGGCCA CGGCTTATGC

50

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-15-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GATCTCTAGA TTAGCTCATT TGTGGTCT

28

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCGGATCCA TGACTGTACT CTACCCA

27

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAC TCGAGGCTCA TTTGTGGTCT

60

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3974 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGTACCTAAG TGAGTAGGGC GTCCGATCGA CGGACGCCTT TTTTTTGAAT TCGTAATCAT

60

GGTCATAGCT GTTTCCTGTG TGAAATTGTT ATCCGCTCAC AATTCCACAC AACATACGAG

120

CCGGAAGCAT AAGTGTAAA CCCTGGGGTG CCTAATGAGT GAGCTAACTC ACATTAATTG

180

CGTTGCGCTC ACTGCCCGCT TTCCAGTCGG GAAACCTGTC GTGCCAGCTG CATTAAATGAA

240

-16-

TCGGCCAACG	CGCGGGGAGA	GGCGGTTTGC	GTATTGGGCG	CTCTTCCGCT	TCCTCGCTCA	300
CTGACTCGCT	GCGCTCGGTC	GTTCCGGCTGC	GGCGAGCGGT	ATCAGCTCAC	TCAAAGGCGG	360
TAATACGGTT	ATCCACAGAA	TCAGGGGATA	ACGCAGGAAA	GAACATGTGA	GCAAAAGGCC	420
AGCAAAAGGC	CAGGAACCGT	AAAAAGGCCG	CGTTGCTGGC	GTTTTTCCAT	AGGCTCCGCC	480
CCCCTGACGA	GCATCACAAA	AATCGACGCT	CAAGTCAGAG	GTGGCGAAAC	CCGACAGGAC	540
TATAAAGATA	CCAGGCGTTT	CCCCCTGGAA	GCTCCCTCGT	GCGCTCTCCT	GTTCCGACCC	600
TGCCGCTTAC	CGGATACCTG	TCCGCCTTTC	TCCCTTCGGG	AAGCGTGGCG	CTTTCTCATA	660
GCTCACGCTG	TAGGTATCTC	AGTTCGGTGT	AGGTCGTTTC	CTCCAAGCTG	GGCTGTGTGC	720
ACGAACCCCC	CGTTCAGCCC	GACCGCTGCG	CCTTATCCGG	TAACATCGT	CTTGAGTCCA	780
ACCCGGTAAG	ACACGACTTA	TCGCCACTGG	CAGCAGCCAC	TGGTAACAGG	ATTAGCAGAG	840
CGAGGTATGT	AGGCGGTGCT	ACAGAGTTCT	TGAAGTGGTG	GCCTAACTAC	GGCTACACTA	900
GAAGAACAGT	ATTTGGTATC	TGCGCTCTGC	TGAAGCCAGT	TACCTTCGGA	AAAAGAGTTG	960
GTAGCTCTTG	ATCCGGCAAA	CAAACCACCG	CTGGTAGCGG	TGGTTTTTTT	GTTTGCAAGC	1020
AGCAGATTAC	GCGCAGAAAA	AAAGGATCTC	AAGAAGATCC	TTTGATCTTT	TCTACGGGGT	1080
CTGACGCTCA	GTGGAACGAA	AACTCACGTT	AAGGGATTTT	GGTCATGAGA	TTATCGTCGA	1140
CAATTCGCGC	GCGAAGGCGA	AGCGGCATGC	ATTACGTTG	ACACCATCGA	ATGGTGCAAA	1200
ACCTTTCGCG	GATATGGCATG	ATAGCGCCCG	GAAGAGAGTC	AATTCAGGGT	GGTGAATGTG	1260
AAACCAGTAA	CGTTATACGA	TGTCGCAGAG	TATGCCGGTG	TCTCTTATCA	GACCGTTTCC	1320
CGCGTGGTGA	ACCAGGCCAG	CCACGTTTCT	GCGAAAAACG	GGGAAAAAGT	GGAAGCGGCG	1380
ATGGCGGAGC	TGAATTACAT	TCCCAACCGC	GTGGCACAAC	AACTGGCGGG	CAAACAGTCG	1440
TTGCTGATTG	GCGTTGCCAC	CTCCAGTCTG	GCCCTGCACG	CGCCGTCGCA	AATTGTGCGG	1500
GCGATTAAAT	CTCGCGCCGA	TCAACTGGGT	GCCAGCGTGG	TGGTGTGAT	GGTAGAACGA	1560
AGCGGCGTCG	AAGCCTGTAA	AGCGGCGGTG	CACAATCTTC	TCGCGCAACG	CGTCAGTGGG	1620
CTGATCATTG	ACTATCCGCT	GGATGACCAG	GATGCCATTG	CTGTGGAAGC	TGCCTGCACT	1680
AATGTTCCGG	CGTTATTTCT	TGATGTCTCT	GACCAGACAC	CCATCAACAG	TATTATTTTC	1740
TCCCATGAAG	ACGGTACGCG	ACTGGGCGTG	GAGCATCTGG	TCGCATTGGG	TCACCAGCAA	1800
ATCGCGCTGT	TAGCGGGCCC	ATTAAAGTCT	GTCTCGGCGC	GTCTGCGTCT	GGCTGGCTGG	1860
CATAAATATC	TCACTCGCAA	TCAAATTCAG	CCGATAGCGG	AACGGGAAGG	CGACTGGAGT	1920

-17-

GCCATGTCCG	GTTTTCAACA	AACCATGCAA	ATGCTGAATG	AGGGCATCGT	TCCCACTGCG	1980
ATGCTGGTTG	CCAACGATCA	GATGGCGCTG	GGCGCAATGC	GCGCCATTAC	CGAGTCCGGG	2040
CTGCGCGTTG	GTGCGGATAT	CTCGGTAGTG	GGATACGACG	ATACCGAAGA	CAGCTCATGT	2100
TATATCCCGC	CGTTAACCAC	CATCAAACAG	GATTTTCGCC	TGCTGGGGCA	AACCAGCGTG	2160
GACCGCTTGC	TGCAACTCTC	TCAGGGCCAG	GCGGTGAAGG	GCAATCAGCT	GTTGCCCCGTC	2220
TCACTGGTGA	AAAGAAAAAC	CACCCTGGCG	CCCAATACGC	AAACCGCCTC	TCCCCGCGCG	2280
TTGGCCGATT	CATTAATGCA	GCTGGCACGA	CAGGTTTCCC	GACTGGAAAG	CGGGCAGTGA	2340
GCGCAACGCA	ATTAATGTAA	GTTAGCGCGA	ATTGTGACCG	AAAGCGGCCA	TCGTGCCTCC	2400
CCACTCCTGC	AGTTCGGGGG	CATGGATGCG	CGGATAGCCG	CTGCTGGTTT	CCTGGATGCC	2460
GACGGATTG	CACTGCCGGT	AGAACTCCGC	GAGGTCGTCC	AGCCTCAGGC	AGCAGCTGAA	2520
CCAACTCGCG	AGGGGATCGA	GCCCCGGGTG	GGCGAAGAAC	TCCAGCATGA	GATCCCCGCG	2580
CTGGAGGATC	ATCCAGCCGG	CGTCCCGGAA	AACGATTCCG	AAGCCCAACC	TTTCATAGAA	2640
GGCGGCGGTG	GAATCGAAAT	CTCGTGATGG	CAGGTTGGGC	GTCGCTTGGT	CGGTCATTTT	2700
GAACCCCAAG	GTCCCGCTCA	GAAGAACTCG	TCAAGAAGGC	GATAGAAGGC	GATGCGCTGC	2760
GAATCGGGAG	CGGCGATACC	GTAAAGCAGC	AGGAAGCGGT	CAGCCCATTG	GCCGCCAAGC	2820
TCTTCAGCAA	TATCACGGGT	AGCCAACGCT	ATGTCCTGAT	AGCGGTCCGC	CACACCCAGC	2880
CGGCCACAGT	CGATGAATCC	AGAAAAGCGG	CCATTTTCCA	CCATGATATT	CGGCAAGCAG	2940
GCATCGCCAT	GGGTCACGAC	GAGATCCTCG	CCGTCGGGCA	TGCGCGCCTT	GAGCCTGGCG	3000
AACAGTTCGG	CTGGCGCGAG	CCCCTGATGC	TCTTCGTCCA	GATCATCCTG	ATCGACAAGA	3060
CCGGCTTCCA	TCCGAGTACG	TGCTCGCTCG	ATGCGATGTT	TCGCTTGGTG	GTCGAATGGG	3120
CAGGTAGCCG	GATCAAGCGT	ATGCAGCCGC	CGCATTGCAT	CAGCCATGAT	GGATACTTTC	3180
TCGGCAGGAG	CAAGGTGAGA	TGACAGGAGA	TCCTGCCCCG	GCACTTCGCC	CAATAGCAGC	3240
CAGTCCCTTC	CCGCTTCAGT	GACAACGTCG	AGCACAGCTG	CGCAAGGAAC	GCCCCTCGTG	3300
GCCAGCCACG	ATAGCCGCGC	TGCCTCGTCC	TGCAGTTCAT	TCAGGGCACC	GGACAGGTGC	3360
GTCTTGACAA	AAAGAACCGG	GCGCCCCCTG	GCTGACAGCC	GGAACACGGC	GGCATCAGAG	3420
CAGCCGATTG	TCTGTTGTGC	CCAGTCATAG	CCGAATAGCC	TCTCCACCCA	AGCGGCCGGA	3480
GAACCTGCGT	GCAATCCATC	TTGTTCAATC	ATGCGAAACG	ATCCTCATCC	TGTCTCTTGA	3540
TCAGATCTTG	ATCCCCTGCG	CCATCAGATC	CTTGGCGGCA	AGAAAGCCAT	CCAGTTTACT	3600

-18-

TTGCAGGGCT TCCCAACCTT ACCAGAGGGC GCCCCAGCTG GCAATTCCGG TTCGCTTGCT 3660
 GTCCATAAAA CCGCCCAGTC TAGCTATCGC CATGTAAGCC CACTGCAAGC TACCTGCTTT 3720
 CTCTTTGCGC TTGCGTTTTT CTTGTCCAG ATAGCCCAGT AGCTGACATT CATCCGGGGT 3780
 CAGCACCGTT TCTGCGGACT GGCTTTCTAC GTGTTCCGCT TCCTTTAGCA GCCCTTGCGC 3840
 CCTGAGTGCT TGCGGCAGCG TGAAGCTTAA AAAACTGCAA AAAATAGTTT GACTTGTGAG 3900
 CGGATAACAA TTAAGATGTA CCCAATTGTG AGCGGATAAC AATTTACAC ATTAAGAGG 3960
 AGAAATTACA TATG 3974

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 112 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AAGCTTAAAA AACTGCAAAA AATAGTTTGA CTTGTGAGCG GATAACAATT AAGATGTACC 60
 CAATTGTGAG CGGATAACAA TTTACACAT TAAAGAGGAG AAATTACATA TG 112

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 419 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met His Ser Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
 1 5 10 15
 Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Phe
 20 25 30
 Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
 35 40 45

-19-

Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 50 55 60
 Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
 65 70 75 80
 Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
 85 90 95
 Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
 100 105 110
 His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
 115 120 125
 Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe
 130 135 140
 Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr
 145 150 155 160
 Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr
 165 170 175
 Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
 180 185 190
 Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
 195 200 205
 Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
 210 215 220
 Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
 225 230 235 240
 Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
 245 250 255
 Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
 260 265 270
 Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
 275 280 285
 Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
 290 295 300
 Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys
 305 310 315 320
 Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu
 325 330 335
 Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro

-20-

340	345	350
Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys		
355	360	365
Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr		
370	375	380
Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser		
385	390	395
Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro		
405	410	415
Gln Met Ser		

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GCAGCACATA TGACAGAAGA GACTATAAAA

30

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GCAGCAGGTA CCTCACAGTT TAGACATGCA

30

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-21-

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GCAGCAGGTA CCTCAACGTC TAATAATGGA

30

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCAGCAGGAT CCCACAGAAG AGACTATAAA

30

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCAGCATCTA GATCACAGTT TAGACATGCA

30

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GCAGCAGGAT CCCACAGAAG AGACTATAAA ATTGGCTGC

39

(2) INFORMATION FOR SEQ ID NO:25:

-22-

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GCAGCATCTA GATCAACGTC TAATAATGGA ATGAAC

36

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 55 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GATCGATCCA TCATGCACTC GCTGGGCTTC TTCTCTGTGG CGTGTTCTCT GCTCG

55

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GCAGGGTACG GATCCTAGAT TAGCTCATTT GTGGTCTTT

39

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

-23-

GACTGGATCC GCCACCATGC ACTCGCTGGG CTTCTTCTC

39

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GACTGGTACC TTATCACATA AAATCTTCCT GAGCC

35

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GACTGGATCC GCCACCATGC ACTCGCTGGG CTTCTTCTC

39

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GACTGGTACC TTATCAGTCT AGTTCTTTGT GGGG

34

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-24-

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GACTGGATCC GCCACCATGC ACTCGCTGGG CTTCTTCTC

39

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GACTGGTACC TCATTACTGT GGACTTTCTG TACATTC

37

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 38 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GCAGCAGGAT CCACAGAAGA GACTATAAAA TTGCTGC

38

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

CGTCGTTCTA GATCACAGTT TAGACATGCA TCGGCAG

37

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/05021

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet
US CL : Please See Extra Sheet
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/4, 6, 69.1, 69.4, 320.1, 325, 243; 436/501; 514/2, 12; 530/387.1, 399

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, EMBASE, WPIDS, BIOSIS
search terms: keratinocyte growth factor (2), kgf, kgf-2, fgf-12, fibroblast growth factor 12

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/06844 A1 (HUMAN GENOME SCIENCES, INC.) 19 February 1998, entire document including claims.	1-19, 22-26

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* B	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L	* A	document member of the same patent family
* O		
* P		

Date of the actual completion of the international search

03 JUNE 1999

Date of mailing of the international search report

11 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHRISTINE SAOUD

Telephone No. (703) 308-0196

JOYCE BRIDGERS
PARALEGAL SPECIALIST
CHEMICAL MATRIX

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/05021

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6): C12N 1/21, 5/10, 15/18, 15/63; C12Q 1/00, 1/68, 1/70; A61K 38/18; C07K 14/50, 16/22

A. CLASSIFICATION OF SUBJECT MATTER:

US CL : 435/4, 6, 69.1, 69.4, 320.1, 325, 243; 436/501; 514/2, 12; 530/387.1, 399